

The Project Gutenberg eBook of The British Woodlice, by Charles Sillem and Wilfred Mark Webb

This ebook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg License included with this ebook or online at www.gutenberg.org. If you are not located in the United States, you'll have to check the laws of the country where you are located before using this eBook.

Title: The British Woodlice

Author: Charles Sillem

Author: Wilfred Mark Webb

Release date: October 10, 2013 [EBook #43928]


Most recently updated: November 1, 2013

Language: English

Credits: Produced by Chris Curnow, Quentin Campbell and the Online

Distributed Proofreading Team at <http://www.pgdp.net> (This file was produced from images generously made available by The Internet Archive)

*** START OF THE PROJECT GUTENBERG EBOOK THE BRITISH WOODLICE ***



Transcriber's Note:

The twenty-five engraved illustrations of adult woodlice shown in Plates I to XXV at the end of the book have been prepared with thumbnails which preserve the same scale as the original book. Clicking on one of these thumbnail images, or on most other figures in the text, will display a larger image to the same scale.

Obvious typographical errors have been corrected. Inconsistent accents, punctuation, and hyphenation are as in the original text.

The "æ" ligature is used interchangeably with "ae" throughout.

Some taxonomic names may have changed since 1906.

For a complete list of corrections, please see the [end of this document](#).

THE BRITISH WOODLICE.

This monograph first appeared in the "Essex Naturalist" (Volume XIV., 1905-6) and has been republished by special arrangement with the Council of the Essex Field Club.

THE BRITISH WOODLICE.

BEING

A MONOGRAPH OF THE TERRESTRIAL ISOPOD
CRUSTACEA

OCCURRING IN THE BRITISH ISLANDS.

BY

WILFRED MARK WEBB, F.L.S.,

*Lecturer on Biology and Nature Study to the Surrey County
Council,*

*Honorary Secretary of the Selborne Society,
Sometime, Senior Assistant Lecturer on Biology to the Essex
County Council,*

*and Editor of the Journal of Malacology,
Joint Author of "Eton Nature Study and Observational
Lessons."*

AND

CHARLES SILLEM.

WITH TWENTY-FIVE PLATES AND FIFTY-NINE FIGURES IN THE TEXT.

LONDON:

DUCKWORTH & CO.,

3, HENRIETTA STREET, COVENT GARDEN.

1906.

PREFACE.

In Professor Sars' "Crustacea of Norway," quite a number

of the British species of woodlice are figured in detail and described in English, but few copies of this fine work are to be met with in our country. The Rev. Canon Norman has from time to time published notes on the British species in "The Annals and Magazine of Natural History;" these are, however, scattered, and contain but few figures, while other literature that exists is out of date. Under these circumstances, we have thought that the following account and figures of all the British species would be useful to those anxious to work at the woodlice, and might also encourage others to pay attention to the distribution and habits of the interesting tribe to which they belong.

The writers would welcome any corrections or additions in view of a second edition.

W.M.W.
C.S.

ODSTOCK, HANWELL, *December, 1905.*

CONTENTS.

[Pg vii]

	PAGE
Introduction	1
Geological history	1
External structure and appendages	2
Alimentary canal	6
Circulatory system	7
Excretory system	7
Nervous system	8
Reproductive organs	8
Development	9
Habits and Economic considerations	12
Local names	15
Methods of collections and preservation	16
Classification	17
Scheme of classification and synopsis of generic characters	18
British Species	19
Section—Ligiæ	19
Family—Ligiidæ	19
Genus— <i>Ligia</i> Fabricius	19
<i>Ligia oceanica</i> Linzè	19
Genus— <i>Ligidium</i> Brandt	21
<i>Ligidium hypnorum</i> Cuvier	21
Family—Trichoniscidæ	22
Genus— <i>Trichoniscus</i> Brandt	22
<i>Trichoniscus pusillus</i> Brandt	22
<i>Trichoniscus vividus</i> Koch	23
<i>Trichoniscus roseus</i> Koch	24
Genus— <i>Trichoniscoides</i> , Sars	25
<i>Trichoniscoides albidus</i> Budde-Lund	25
Genus— <i>Haplophthalmus</i> Schobl	26
<i>Haplophthalmus mengii</i> Zaddach	26
<i>Haplophthalmus danicus</i> Budde-Lund	27
Family—Oniscidæ	27
Genus— <i>Oniscus</i> Linné	27
<i>Oniscus asellus</i> Linné	27
Genus— <i>Philoscia</i> Latreille	29
<i>Philoscia muscorum</i> Scopoli	29
<i>Philoscia couchii</i> Kinahan	30
Genus— <i>Platyarthrus</i> Brandt	30
<i>Platyarthrus hoffmannseggii</i> Brandt	30
Genus— <i>Porcellio</i> Latreille	32
<i>Porcellio scaber</i> Latreille	32
<i>Porcellio pictus</i> Brandt and Ratzeburg	33
<i>Porcellio dilatatus</i> Brandt	33
<i>Porcellio rathkei</i> Brandt	34
<i>Porcellio laevis</i> Latreille	35
<i>Porcellio ratzeburgii</i> Brandt	36
Genus— <i>Metoponorthus</i> Budde-Lund	37
<i>Metoponorthus pruinosus</i> Brandt	37
<i>Metoponorthus cingendus</i> Kinahan	38

[Pg viii]

Genus—Cylisticus Schnitzler	38
<i>Cylisticus convexus</i> De Geer	39
Family—Armadillidiidæ	40
Genus—Armadillidium Brandt	40
<i>Armadillidium nasatum</i> Budde-Lund	40
<i>Armadillidium vulgare</i> Latreille	41
<i>Armadillidium pulchellum</i> Zencker	42
<i>Armadillidium depressum</i> Brandt	43
Distribution of species	43
Conclusion	43
Bibliography	45

LIST OF FIGURES.

[Pg ix]

PLATES I.-XXV. with a List, will be found at the end of the Book.

FIGURE.		PAGE.
1.	Parts of the body (<i>Oniscus asellus</i>)	2
2.	The first antenna (<i>Oniscus asellus</i>)	2
3.	The second antenna (<i>Oniscus asellus</i>)	3
4.	The underside of the head (<i>Oniscus asellus</i>)	3
5.	The mandibles (<i>Oniscus asellus</i>)	4
6.	The first maxillae (<i>Oniscus asellus</i>)	4
7.	The second maxillae (<i>Oniscus asellus</i>)	4
8.	The fused maxillipeds (<i>Oniscus asellus</i>)	4
9.	The "upper lip" (<i>Oniscus asellus</i>)	5
10.	The "lower lip" (<i>Oniscus asellus</i>)	5
11.	A typical thoracic segment (<i>Oniscus asellus</i>)	5
12.	The fifth thoracic segment of a female (<i>Oniscus asellus</i>)	5
13.	The underside of the abdomen of a female (<i>Oniscus asellus</i>)	6
14.	A typical abdominal appendage (<i>Oniscus asellus</i>)	6
15.	The first abdominal appendage of the male (<i>Oniscus asellus</i>)	6
16.	The second abdominal appendage of the male (<i>Oniscus asellus</i>)	6
17.	The alimentary canal (<i>Oniscus asellus</i>)	7
18.	The circulatory system (<i>Oniscus asellus</i>)	7
19.	The nervous system (<i>Oniscus asellus</i>)	8
20.	Female reproductive organs (<i>Oniscus asellus</i>)	8
21.	The male reproductive organs (<i>Oniscus asellus</i>)	9
22.	The fertilized egg (<i>Porcellio scaber</i>) after Roule	10
23.	The fertilized egg seen in section (<i>Porcellio scaber</i>) after Roule	10
24 to	The development of a woodlouse (<i>Porcellio scaber</i>) after	10
31.	Roule	
32.	Embryo of the woodlouse showing the three divisions of the intestine separately developed (<i>Porcellio scaber</i>) after Roule	11
33.	Embryo of the woodlouse showing traces of the segments (<i>Porcellio scaber</i>) after Roule	11
34.	An embryo woodlouse ready to be hatched (<i>Porcellio scaber</i>) after Roule	11
35.	Flagellum and last peduncular joint of the antenna of <i>Ligia oceanica</i>	12
36.	Flagellum and last peduncular joint of the antenna of <i>Ligidium hypnorum</i>	13
37.	Flagellum and last peduncular joint of the antenna of <i>Trichoniscus pusillus</i>	23
38.	Flagellum and last peduncular joint of the antenna of <i>Trichoniscus vividus</i>	24
39.	Flagellum and last peduncular joint of the antenna of <i>Trichoniscus roseus</i>	24
40.	Flagellum and last peduncular joint of the antenna of <i>Trichoniscoides albidus</i>	25
41.	Flagellum and last peduncular joint of the antenna of <i>Haplophthalmus mengii</i>	26
42.	Flagellum and last peduncular joint of the antenna of <i>Haplophthalmus danicus</i>	27
43.	Flagellum and last peduncular joint of the antenna of <i>Oniscus asellus</i>	28
44.	Flagellum and last peduncular joint of the antenna of <i>Philoscia muscorum</i>	29
45.	Flagellum and last peduncular joint of the antenna of <i>Philoscia couchii</i>	30
	Flagellum and last peduncular joint of the antenna of	

[Pg x]

46.	<i>Platyarthrus hoffmannseggii</i>	31
47.	Flagellum and last peduncular joint of the antenna of <i>Porcellio scaber</i>	32
48.	Flagellum and last peduncular joint of the antenna of <i>Porcellio pictus</i>	33
49.	Flagellum and last peduncular joint of the antenna of <i>Porcellio dilatatus</i>	34
50.	Flagellum and last peduncular joint of the antenna of <i>Porcellio rathkei</i>	35
51.	Flagellum and last peduncular joint of the antenna of <i>Porcellio laevis</i>	35
52.	Flagellum and last peduncular joint of the antenna of <i>Porcellio ratzeburgii</i>	36
53.	Flagellum and last peduncular joint of the antenna of <i>Metoponorthus pruinosus</i>	37
54.	Flagellum and last peduncular joint of the antenna of <i>Metoponorthus cingendus</i>	38
55.	Flagellum and last peduncular joint of the antenna of <i>Cylisticus convexus</i>	39
56.	Flagellum and last peduncular joint of the antenna of <i>Armadillidium nasatum</i>	40
57.	Flagellum and last peduncular joint of the antenna of <i>Armadillidium vulgare</i>	41
58.	Flagellum and last peduncular joint of the antenna of <i>Armadillidium pulchellum</i>	42
59.	Flagellum and last peduncular joint of the antenna of <i>Armadillidium depressum</i>	43

THE BRITISH WOODLICE.

[Pg 1]

Introduction.—Having finished a somewhat exhaustive list of the land and fresh-water molluscs of Essex, ^[1] one of the present writers felt that if he were to make any further contributions of importance to a knowledge of the fauna of that interesting county, he must turn his attention to some other group of animals. It seemed most fitting that some creatures should be chosen which are commonly met with during the search for molluscs. Centipedes, millepedes, and woodlice fulfilled these conditions, and all were collected, but as only seventeen species of woodlice had at the time been found in England, it was deemed advisable to study these in detail to begin with. The present contribution is the result of the undertaking, and we have thought that a general consideration of the British Woodlice, with careful drawings from nature of all the species now known from this country, ought to lead to a more general study of these interesting creatures and their habits.

Position in the scheme of classification.—The Woodlice belong to an immense group of invertebrate animals known as the Arthropoda, the bodies of which are segmented and provided with jointed appendages for purposes of walking, swimming, and feeding. Of this group, two large divisions are recognized. The first contains the forms which breathe by means of air-tubes, such as the Insects; and the second has been constituted for Crustacea, which breathe by means of gills. The latter are, of course, adapted more especially for a life in water, but here and there we come across examples so modified that they can exist in air. The land-crabs are a case in point, and so are the Woodlice. These belong to an order which contains many fresh-water and marine species, known as the Isopoda.

Geological history.—The known history of the order is a long one, for remains occur in the Old Red Sandstone (Devonian) of Herefordshire, and in the Coal Measures. (79) ^[2] A form which has been named *Archæoniscus brodiei*, and is said to be referable to the recent family Aegidae which is found in some numbers in the Purbeck Beds (Upper Jurassic), of this country (47). Fossil Isopods have also been recorded from the Oolite and from the Oligocene (Isle of Wight).

[Pg 2]




FIG. 1.—PARTS OF THE BODY.
(*Oniscus asellus*.)

Turning to the Woodlice proper, we find that they first make their appearance in the Miocene (of Oenigen and Baden), and occur also in amber (79); while examples of genera, such as *Oniscus* and

Porcellio, have been discovered in late Tertiary deposits (47).

External structure and appendages.

—Woodlice agree in being of a somewhat oval form, and their bodies are arched, the curve varying in different genera and species. A *head* is to be distinguished; behind this comes the *thorax* of seven segments which are often considerably broader than the six succeeding ones which form the *abdomen* (see fig. 1.)

The head carries two *large antennae* (fig. 3) which are very evident, and a careful search with a lens will reveal a second and minute pair (the *smaller antennae*) situated between the base of the others, and really anterior to them. (figs. 2 and 4.)

The larger antennae are customarily bent at certain points, and we can distinguish a terminal part, or *flagellum*, and a basal part, the *peduncle* (fig. 3). The number of joints in these structures, which varies in different genera and species, forms a useful classificatory character, and the relative length of the component parts is of considerable value in distinguishing species.




FIG. 3.—THE SECOND ANTENNA.
(*Oniscus asellus*.)

There are four pairs of mouth




FIG. 4.—THE UNDERSIDE OF THE HEAD.
(*Oniscus asellus*.)

appendages—namely the jaws or *mandibles* (fig. 5), the *first maxillae* (fig. 6), the *second maxilla* (fig. 7), and the *maxillipeds* (fig. 8). When the head is examined from the underside the last of these organs will be seen first, covering in the others.

A small median plate attached to the front of the head has been called "*the upper lip*" (fig. 9), while inside the mouth appendages is a little bilobed structure "*the lower lip*" (fig. 10).

Before leaving the external features of the head, we must allude to the pair of *eyes* which are usually present, though never raised on stalks. In the Common Woodlouse (*Oniscus asellus*, from which all our figures to illustrate structure have been made), as in many other species, the eyes are compound (fig. 4), but in some forms these are simple.




FIG. 5.—THE MANDIBLES.
(*Oniscus asellus*.)




FIG. 6.—THE FIRST MAXILLAE.
(*Oniscus asellus*.)




FIG. 7.—THE SECOND
MAXILLAE.
(*Oniscus asellus.*)




FIG. 8.—THE FUSED
MAXILLIPEDS.
(*Oniscus asellus.*)

Each of the seven joints of the thorax bears a pair of *walking legs* (fig. 11), and in the female at the time when the eggs are laid, a pair of plates (fig. 12) arises on segments II. to V. These plates together form a brood pouch, in which the eggs are carried (fig. 12) until they are hatched, and in which the young ones remain for some time afterwards.

[Pg 5]



FIG. 9.—THE
"UPPER LIP."
(*Oniscus asellus.*)



FIG. 10.—THE
"LOWER LIP."
(*Oniscus asellus.*)




FIG. 11.—A TYPICAL THORACIC SEGMENT.
(*Oniscus asellus.*)




FIG. 12.—THE FIFTH THORACIC SEGMENT OF A FEMALE.
(*Oniscus asellus.*)

When we examine the abdomen, we find that the appendages are plate-like, with the exception of the last pair (fig. 13), and they all agree in having two divisions, an arrangement which would prove awkward in limbs used for walking or feeling.




FIG. 13.—THE UNDERSIDE OF THE
ABDOMEN OF A FEMALE.
(*Oniscus asellus.*)

The inner plate (or endopodite) is in structure a *gill*, but the blood that passes through it, is enabled to take up oxygen from moist air, while the outer division (or exopodite) acts as a protecting cover (fig. 14). In *Porcellio*, air-tubes (*tracheae*) may be




FIG. 15.—THE FIRST
ABDOMINAL APPENDAGE
OF THE MALE.
(*Oniscus asellus.*)

[Pg 6]

present (see below).

In the male, the first two pairs of abdominal appendages are specially modified, the inner divisions (endopodites) being long and pointed (figs. 15 and 16). The last pair, or tail appendages, in the male are often considerably larger than in the female, and the form of these structures is sometimes of value in classification.




FIG. 14.—A TYPICAL ABDOMINAL APPENDAGE. (*Oniscus asellus*.)

Alimentary canal.

—The main portion of the alimentary system is, practically speaking, a straight tube (fig. 17). Its first part (not shown in the figure) is a narrow gullet,

which after passing through the nerve collar dilates to form a sort of stomach. Into this the secretion of four digestive glands is poured by two ducts. These glands have a somewhat striking appearance, being yellow tubes spirally coiled, and they end blindly. From the stomach the intestine runs to the hinder end of the body and passes under the heart.




FIG. 16.—THE SECOND ABDOMINAL APPENDAGE OF THE MALE. (*Oniscus asellus*.)

Circulatory system.—The blood being aerated in the abdominal appendages, we find that the heart is situated towards the hinder end of the body (fig. 18). Three main arteries supply the thorax and head, while the blood is brought from the gills to the heart.

[Pg 7]




Fig. 17.—THE ALIMENTARY CANAL. (*Oniscus asellus*.)

Excretory system.

—The excretory organs consist of a (a) pair of so-called "shell glands," which are considered to be the equivalents of the excretory tubes or nephridia of annelid worms. In the woodlouse these excretory organs open on the second pair of maxillae. They are composed of a tube (*sacculus*) closed at one end and more or less bent upon itself (5,




Fig. 18.—THE CIRCULATORY SYSTEM. (*Oniscus asellus*.)

p. 261) which communicates with a *labyrinth* that is provided with an excretory orifice. Matters are eliminated by the *epithelial cells* [the histology has been described and figured in *Ligidium hypnorum* (66)], which are very large in *Ligia oceanica*.

(b) Masses of cellules in the head, very greatly developed in *Ligia oceanica* (but numbering scarcely more than ten in *Oniscus asellus*), which have no external opening. They also function as excretory organs (5, p. 263), and have been called "cephalic nephrocytes."

[Pg 8]

(c) Other "branchial nephrocytes" are situated on the dorsal surface between the last thoracic and the first abdominal segments, as well as between those that follow, with the exception of the last two; they are in distinct patches, one on each of the middle line in *Ligia*, but more or less continuous in *Oniscus* (5, p. 265).

(d) The digestive glands have also been shown to be excretory (5, p. 270).




FIG. 19.—THE NERVOUS SYSTEM.
(*Oniscus asellus*.)




FIG. 20.—FEMALE REPRODUCTORY
ORGANS.
(*Oniscus asellus*.)

Nervous system.—The nervous system consists of *paired ganglia* in the head, above the alimentary canal which send off nerves (*commissures*) that meet below, to form a *double nerve cord* with ganglia at intervals (see fig. 19).

Reproductive organs.—In the female there are a pair of *ovaries* in the positions shewn in fig. 20; and *ducts* run to the underside of the fifth thoracic segment.

The openings are very difficult to identify, and Lereboullet (39, p. 113) was unable to find them. It is obvious that the openings must be underneath the plates that form the egg pouch, and as a change of skin is required to set these free, it would appear that at ordinary seasons the ducts from the ovaries are closed.

[Pg 9]




FIG. 21.—THE MALE
REPRODUCTORY ORGANS.
(*Oniscus asellus*.)

The writers have been able to determine from external examination of specimens which had moulted and were about to lay eggs, that the oviducts at such time open to the inside of the base of each walking leg on the fifth segment. In similar specimens the oviducts were also followed to the opening from within. The *brood pouch* has already been described.

The male organs consist of six *testes* arranged in two pairs, each of which is provided with a *reservoir* (see fig. 21). The efferent ducts from the two reservoirs unite at the base of the thorax to form a common duct (or "penis").

Development.—The eggs, in the common species of woodlice, at least, are laid at the beginning of summer, and are retained in the brood pouch, where they undergo their development. The process has been recently traced with great care by Professor Louis Roule (58) in *Porcellio scaber* and the description which follows is based upon his researches.

As, practically speaking, the larval stages are passed within the egg, and there is no free embryo differing in form from the parent, it is necessary for the young creatures to be well supplied with nutritive material. In fact, the bulk of the large egg is made up of *food-yolk*, on the outside of which the *formative protoplasm* is disposed in irregular patches. In the fertilized ovum, one of the latter, which lies in a particular position at the end, is found to be larger than the others (see fig. 22). It contains the nucleus of the egg-cell (see fig. 23) and is called the *cicatricula*. This is the only portion of the egg which divides and produces *nucleated cells*. It is these which gradually spread all over the surface of the food-yolk, forming a layer known as the *blastoderm*, which is at first but one cell thick (see figs. 24, 26, and 28).

[Pg 10]

Before, however, the food-yolk is quite closed in, a differentiation into two layers—the *pro-ectoderm* and *pro-endoderm*—takes place (see fig. 25) and rudiments of the first two pairs of *appendages* appear (see fig. 26). Moreover, the cells of the ectoderm change their shape and begin to multiply at two points to form the beginnings of the cerebral ganglia and the nerve cord respectively.




FIG. 22.—THE FERTILIZED EGG
(*Porcellio scaber*), AFTER ROULE.




FIG. 23.—THE FERTILIZED EGG SEEN IN SECTION
(*Porcellio scaber*), AFTER ROULE.

As the blastoderm closes over the food-yolk, two more appendages arise and these are soon followed by others (see fig. 28). A depression appears at the point where the blastoderm closed and internally the pro-endoderm or inner layer is differentiated into two—the *endoderm proper* and the *mesoderm* (see fig. 29). The former begins to grow so that its edges unite to form the middle part of the intestine (see fig. 29) seen from the outside in fig. 30. The depression already mentioned grows deeper, forming a tube which is the hind portion of the *intestine*, while at the anterior end of the embryo the front part of the intestine is similarly formed (see fig. 30). By this time also all the nineteen appendages have made their appearance and the mesoderm, (which has grown considerably, to form the beginnings of the muscles) has sent prolongations into each of them. About this time, spaces (see fig. 31) are formed in the muscular mesoderm which are all that remain of the *true body cavity* characteristic of animals above the level of the jelly fish, and in these spaces the blood ultimately circulates.

SURFACE VIEWS.

OPTICAL SECTIONS.



FIG. 24.




FIG. 25.




FIG. 26.




FIG. 27.




FIG. 28.




FIG. 29.




FIG. 30.




FIG. 31.

THE DEVELOPMENT OF A WOODLOUSE (*Porcellio scaber*), AFTER ROULE.

Figs. 24, 26, 28, 30, are Surface Views, and figs. 25, 27, 29, 31, which indicate slightly later stages respectively than the others, are of egg seen in Optical Section.

The body next alters somewhat in shape and the three divisions of the intestine approach one another (see fig. 32) previous to their junction. As may be imagined during these processes the food-yolk has gradually been used up and the space which it occupied taken by the internal organs, which we have mentioned.

[Pg 11]




FIG. 32.—EMBRYO OF THE WOODLOUSE SHOWING THE THREE DIVISIONS OF THE INTESTINE SEPARATELY DEVELOPED (*Porcellio scaber*), AFTER ROULE.




FIG. 33.—EMBRYO OF THE WOODLOUSE SHOWING TRACES OF THE SEGMENTS (*Porcellio scaber*), AFTER ROULE.

In the last stages of the development the appendages become larger still, the *heart* makes its appearance, segmentation of the body is completed, and except that the seventh pair of walking legs are as yet rudimentary the woodlouse is completed. It is only after hatching that the pair of legs mentioned, attain to their normal length.




FIG. 34.—EMBRYO OF THE WOODLOUSE SHOWING TRACES OF THE SEGMENTS (*Porcellio scaber*), AFTER ROULE.

The process of segmentation of the egg and the formation of its layers lasts about a fortnight, while the completion of the development proceeds much more rapidly, for another three weeks bring it to an end.

After the first moult or change of skin the last pair of walking legs makes its appearance, and Mr. James B. Casserley [whose work one of us (75) has described elsewhere] found when keeping a number of the common pill-woodlouse (*Armadillidium vulgare*) in captivity that his specimens did not subsequently change their skins more than once in the six months during which he had them under observation. He also noted that the crustaceans go on growing after they are sexually mature. As his specimens grew older, Mr. Casserley noticed that their colour became darker, and a curious point recorded by him is that two examples of the same age may change their skins at the same time, and while one may have afterwards nearly twice as many markings, on the other very few at all will be seen. The time required for the growth of a woodlouse from the size of a pin's head to that of an adult example—say three-quarters-of-an-inch long—must be fairly considerable, taking into account the fact that any appreciable increase in size can only occur at a moult and Mr. Casserley's observations as to the infrequency of the process in *Armadillidium vulgare*. ([See p. 13.](#))

Habits and Economic Considerations.—The construction of the breathing organs of woodlice, and the necessity which exists for these to be kept moist, restricts the habitats of the animals considerably. Woodlice are found under stones and logs, beneath the bark of dead and rotten trees, among decaying vegetable matter as well as living grass and moss in damp or wet situations. When looking for some of the common species under the bark of fallen trees it is surprising to notice that the crustaceans may be entirely absent from many trunks, while when another is examined which seems to differ very slightly, if at all, in condition or situation, they are found in swarms. There is no doubt but that the habits of woodlice would well repay the attention of naturalists, who are now recognizing that besides anatomy as such, and the classification which a knowledge of structure permits, there is the equally important consideration of the creatures as they live their own life and affect that of others. It is not our object to give a detailed account of the ecology of British woodlice, but rather to provide a basis from which it may be approached. Nevertheless a few general remarks may not come amiss. Many points in the life-history of woodlice may no doubt be learned by keeping them in captivity and there is just sufficient difficulty in doing this successfully to give an interest to the matter.

Apart from a supply of proper food, we take it that the chief object to be attained is the provision of the amount of moisture required by the particular species under examination, together with a sufficient supply of air.

A great many interesting observations can be thus carried out, such as those of Mr. Casserley, to which allusion has already been made. The process of moulting for instance is well worth watching, and although specimens with half their coat changed may be found in remote corners, yet the whole course of the moult can be seen much better in the case of captive woodlice. The following account is taken from Mr. Casserley's description (75) of what happens in the case of *Armadillidium vulgare*:—The approach of the moult is indicated by the appearance of a white border on each segment of the body, which becomes gradually more marked, while at the same time the animal is seen to be less active and often makes a small burrow in which to hide. Sometimes a sheltered corner against a stone is looked upon as affording sufficient protection, but in either case each woodlouse keeps to the place originally chosen. About ten days after the white lines have become visible the animal appears to be divided into two. Its skin is becoming loose and little movement can take place at the joints of its body with the exception of that between the fourth and fifth thoracic segments where the skin will ultimately break. The woodlouse spends a day or two in this condition and then, by suddenly walking forward, frees itself from the covering of

the hinder portion of its body. The three last pairs of walking legs are carefully pulled out from the old skin, which now appears perfectly white, and at the same time the lining of the hind portion of the alimentary canal (hind gut) is also shed. After putting the tender half of his body well into his corner or burrow the woodlouse proceeds to eat the part of his skin that he has cast. The creature has now a very odd appearance. His front half with the exception of the white edges is as it was before, the rest of him instead of a light slaty blue, and is very soft as well as proportionately a little larger.

In three days or so the tail end becomes hard and attains the normal colour. Then the old skin from the front half is pushed off and the creature becomes practically defenceless, so much so in fact, that any of his species that happen to find him will attack him and eat all his front half, rejecting, however, his now hardened tail-end.

[Pg 14]

Provided that the moulting woodlouse has survived (and in captivity, to ensure this, he must be isolated), after three days his jaws will be sufficiently hardened to allow of his eating, and usually he first of all devours the second half of his cast skin. The operation of moulting does not occupy quite so long a time in the case of young examples. Specimens half-an-inch long do not moult more than once in six months and show but little increase in size after the process.

Woodlice do not appear to live on either animal or vegetable food alone, but adopt a mixed diet. It is, however, owing to their attacks upon cultivated plants that the creatures are looked upon as pests by the horticulturalist. The animals feed either in the night or in the very early morning, on seedlings, orchid tubers, mushrooms, or anything that comes to hand. Few of the accounts, however, of their ravages, mention that the crustaceans have been caught absolutely in the act of doing the damage ascribed to them. Some careful inquiries have nevertheless enabled us to discover several observers who have watched woodlice feeding. Mr. F. V. Theobald, of Wye College, and one of the students at Swanley Horticultural College are among the number. The former has also given us an account of the methods, out of many tried, which he has found most successful for getting rid of the crustaceans. Out of doors trapping with moss, sacking or horse-dung is best. In glass houses, fumigation with hydro-cyanic acid gas has cleared them out, and poison baits, especially potatoes cut and soaked in white arsenic, have done some good. Stable manure is especially favourable to these creatures, particularly when it is used "long": in this condition it should therefore be avoided.

It is interesting to note how the woodlice in winter simply remain where they happen to be so long as there is sufficient moisture, though they are ready to run about as rapidly, for a time, as in summer, should they happen to be disturbed.

No doubt many points of inter-relation between woodlice and other animals remain to be discovered. Mr. John W. Odell tells us that on Exmoor, in the open, he found no *Armadillidia*, though other forms occurred under nine out of every ten stones that he turned over, and here the smaller species of ants also abounded. Close to stone walls *Armadillidia* were to be seen to the exclusion of all other genera, and this state of affairs was ascribed by Mr. Odell to the presence of swarms of the large wood-ants which he considers would make short work of any woodlice that could not protect themselves by rolling up.

[Pg 15]

We ought not to conclude this account without mentioning the fact that woodlice once played an important part in medicine.

Doctor Fernie (28) gives some interesting extracts with regard to the hog-louse and the woodlouse. The latter he seems to have identified quite correctly as *Oniscus asellus*. He calls the former, however, indiscriminately, "the common armadillo" (which is the old name for the pill-woodlice now

known as *Armadillidium*), "the pill millipede" and "*Glomeris marginata*." The last two names are those of another creature, not a crustacean, which when it is rolled up can be very easily mistaken for an *Armadillidium*, though, when it uncurls, it will be seen to have many more than seven pairs of legs. The local appellations applied to the hog-louse by Doctor Fernie, and his remarks with regard to its commonness, tend to show that it is *Armadillidium vulgare*, to which he really refers, and the use of which in medicine was commonly general.

Hog-lice were prescribed for scrofulous diseases and obstructions of the liver and digestive organs, among other things, and the London College of Physicians directed that the creatures should be prepared by suspending them in a thin canvas bag placed within a covered vessel over the steam of hot spirit or wine, so that being killed by the spirit they might become friable. Hog-lice and Woodlice were also administered alive, while the former were also put down the throats of cows "to promote the restoration" of their cud, hence their name of "cud-worm." There seems to be considerable evidence that even in modern times Woodlice have had considerable remedial effect which depends upon "an alkaliescent fluid" contained in them.

Local Names.—Among the local names by which these creatures are known are those of "sow bug," "lucre pig" (Berkshire), "carpenter" and "chiselhog" (Berkshire). Doctor Fernie (28) gives a number of others:—"thrush-louse," "tiggyhog," "cheslip," "kitchenball," "chiselbob," "lugdor," "palmer," and "cudworm." In the eastern counties the same writer notes that they are known as "old-sows" or "St. Anthony's hogs" while the Welsh call them "little grey-hogs," "the little old women of the wood" or "grammar-sows," grammar signifying a shrivelled up old dame. *Oniscus asellus* was sometimes called "socchetre," "church louse," and "chinch."

[Pg 16]

Methods of Collection and Preservation.—Woodlice should be collected straightway into tubes or bottles half filled with 30 per cent. methylated spirit.^[3] Woodlice dropped into this weak spirit become gradually narcotised and die, and they remain limp enough for purposes of examination or to allow, of their legs and antennæ being set out during the process of mounting. Specimens to be kept permanently should be placed in 70 per cent. alcohol. For storage purposes the specimens of each species from a given locality should be put together into a small flat bottomed tube such as is used for pillules by apothecaries or specially made for natural history purposes. A paper label on which the name, locality, date of capture and any other necessary particulars have been written with dark lead pencil, is not affected by the spirit. The tubes may be corked, though if not frequently examined all the spirit may evaporate, and cause the specimens to be spoilt. A safer method is to plug the tubes with cotton wool and keep all those containing a given species or specimens from a particular locality beneath the surface of spirit in a large wide-mouthed bottle, into which first of all some cotton wool has been put to prevent the tubes from coming into sudden contact with the glass at the bottom. For show purposes in museums, specimens taken direct from 30 per cent. spirit should be mounted on slips of opal glass by means of gum-tragacanth which has been powdered and shaken up in spirit before having water added to it. The slips can be exhibited in glass tubes, six inches high by one across, or in narrow stoppered museum jars. A variation of the method is to mount the animals on clear glass and to place behind them another strip of any colour that may be preferred.

Classification.—The various genera of woodlice are connected together so closely, by intermediate forms, that their division into families is, to a very great extent, arbitrary. Bate and Westwood described but a single family Oniscidæ (I), though they distinguished two sub-families:—Ligiinæ, which included the forms with many joints to the

[Pg 17]

flagellum of the antenna, and Oniscinæ, which contained the rest.

Since then the pill-woodlice have been thought by some to be sufficiently different from the other genera to warrant their separation, and three families namely, Ligiidæ, Oniscidæ, and Armadillidæ have been recognized, as for instance by Dr. Scharff (63).

A fourth family—Trichoniscidæ—has been added by Professor G. O. Sars, who in his *Crustacea of Norway* (59) alludes to the division of the tribe into the sections Ligie and Onisci and has adopted the following classification:—

Order—**ISOPODA.**

Tribe—ONISCOIDA.

Family I.—LIGIIDÆ.

Ligia.
Ligidium.

Family ONISCIDÆ.
III.—

Oniscus.
Philoscia.
Platyarthrus.
Porcellio.
Metoponorthus.
Cylisticus.

Family II.—TRICHONISCIDÆ.

Trichoniscus.
Trichoniscoides.
Haplophthalmus.

Family ARMADILLIDIÆ.
IV.—

Armadillidium.

All the genera described by Professor Sars are represented in the British Islands.

Below is a scheme of classification and synopsis of the characters of British genera of woodlice which we have compiled in order to render easy the determination of the genus to which any particular specimen may belong.

*SCHEME OF CLASSIFICATION AND SYNOPSIS OF
GENERIC CHARACTERS.*

[Pg 18]

Order—**ISOPODA.**

Tribe—ONISCOIDA.

Section I.—**LIGIÆ.**

The Two Divisions of the Tail Appendages alike in Shape.

(A.)—Flagellum with 10 or more joints;
tail appendages

wholly visible; head without lateral
lobes

LIGIIDÆ.

(1.)—Abdomen broad; body
large; habitat,

the sea-shore

Ligia.

(2.)—Abdomen narrow; habitat,
wet moss

Ligidium.

(B.)—Flagellum with less than 10 joints;
head with

small lateral lobes, tail appendages
partly

covered

TRICHONISCIDÆ.

(3.)—Abdomen narrow; eyes
compound;

flagellum usually with
more than 3

joints

Trichoniscus.

(4.)—Abdomen narrow; eyes
simple or wanting;

flagellum with 4 joints *Trichoniscoides*.
 (5.)—Abdomen broad
 (comparatively); eyes
 simple; back with
 longitudinal
 ridges; flagellum with 3
 joints *Haplophthalmus*.

Section II.—ONISCI.

The Outer Divisions of the Tail Appendages Broader than the Inner.

(A.)—Tail appendages projecting when the animal is
 walking *ONISCIDÆ*.
 (a.)—Unable to roll up into a complete ball.
 (6.)—Flagellum with 3 joints;
 abdomen broad; head, with lateral lobes *Oniscus*.
 (7.)—Flagellum with 3 joints;
 abdomen narrow; head without lateral lobes *Philoscia*.
 (8.)—Flagellum with 1 joint; eyes wanting; abdomen broad; habitat, ant's nests *Platyarthrus*.
 (9.)—Flagellum with 2 joints;
 abdomen broad; frontal lobe projecting *Porcellio*.
 (10.)—Flagellum with 2 joints;
 abdomen narrow *Metoponorthus*.
 (b.)—Able to roll up into a complete ball.
 (11.)—Flagellum with 2 joints; antennae folded together over the thorax when the animal is rolled up into a ball *Cylisticus*.
 (B.)—Tail appendages not projecting when the animal
 walking *ARMADILLIDIIDÆ*.
 (12.)—Flagellum with 2 joints;
 antennae hidden or carried at the sides of the head when the animal is rolled up into a ball *Armadillidium*.

British Species.—Naturalists in this country paid little attention to the recognition or description of Woodlice, until the latter half of the nineteenth century.

In 1857 Kinahan read a paper before the British Association (32) in which he described fourteen species of woodlice from the British Islands, and eleven years later when Bate and Westwood published their book (1), the number had risen to seventeen. One of the species (*Oniscus fossor*), however, was doubtful, and although Dr. Scharff in 1894 (63) rejected it, his list contained also seventeen species, for in the meantime the Rev. T. R. R. Stebbing had found *Ligidium hypnorum* in Surrey (70).

Since then the Rev. Canon Norman, Dr. Scharff, the Rev. T. R. R. Stebbing, and one of the present writers, have added other species, as will be seen from the following pages, in

which all those found, up to the present time in the British Islands are described and figured.

We shall now consider in detail the British genera and species of woodlice and give their synonymy and distribution.

Order—**ISOPODA.**

Tribe—**ONISCOIDA.**

Section—**LIGIÆ.**

THE TWO DIVISIONS OF THE TAIL APPENDAGES ALIKE
IN SHAPE.

Family—**LIGIIDÆ.**

**Flagellum with ten or more joints; tail appendages
wholly visible; head without lateral lobes.**

Genus—**LIGIA** Fabricius, 1798 (27), p. 301.

Abdomen broad; body large; habitat, the sea-shore.

The genus *Ligia* agrees with *Ligidium* alone, in that the flagellum of the larger antennæ has more than ten joints. In both genera, there are no lateral lobes to the head, and the tail appendages are wholly visible from the upper surface of the body. The latter in *Ligia* is, however, very many times bigger than in *Ligidium* and shows no abrupt decrease in the width of its segments when the abdomen is reached.

Ligia oceanica Linné (The Quay-louse). [PLATE I.](#)

[Pg 20]

1767 *Oniscus oceanicus* Linné (43), p. 1061.

1793 *Cymothoa oceanica* Fabricius (26), p. 509.

1815 *Ligia scopulorum* Leach (38), p. 374.

1868 *Ligia oceanica* Bate and Westwood (1), p. 444.

1898 *Ligia oceanica* Sars (59), 156, pl. LXX.




FIG. 35.—FLAGELLUM
AND LAST PEDUNCULAR
JOINT OF THE ANTENNA
OF *Ligia oceanica*.

There is but one British species of *Ligia*, and this, the largest member of the whole tribe to be met with in these Islands, usually attains a length of two centimetres, while adult males may be nearly half as long again. It is the *Oniscus oceanicus* of Linnæus and lives on the sea shore, where it may be found at low tide beneath stones and rubbish in the crevices of timber. *Ligia* forms a connecting link between the woodlice proper and the many Isopods which actually live in the sea.

The colour of the animals is a greenish grey, and the compound eyes are almost black, so that they are very conspicuous; there are from eleven to fourteen joints to the flagellum of the outer antennæ and this feature, taken in conjunction with the large size and habitat, is sufficient to identify the species in question.

On the coast of Essex the name "quay-lowders" is given to these crustaceans, "lowder" being apparently an old plural of louse.

It is worthy of mention that Mr. Webb, when in charge of the Marine Biological Station at Brightlingsea, examined a very large male specimen of *Ligia oceanica*, in which the maxillæ were duplicated and consisted of four pairs instead of two.

BRITISH LOCALITIES:—

England: Brightlingsea; (W.M.W.): Maldon; (W.M.W. from R.M.): Southend; (J.A.M.): Whitstable; (W.M.W.): Herne Bay; Margate; Dover; Folkestone; (J.A.M.)

Scotland: Shetland to Cornwall; (Norman, 49).

Ireland: East Coast; West Glengariff; Castletown; Berehaven; Bundoran; (Scharff, 63).

FOREIGN DISTRIBUTION:—

Europe: France; (25); Spain; (12); Denmark; Prussia; Norway; Faroe Islands; Belgium; (59).

Africa: Morocco; (16).

[Pg 21]

Genus—**LIGIDIUM** Brandt, 1833 (3), p. 173. Zia, Koch (34).

Abdomen narrow; habitat, wet moss.

In *Ligidium* there are numerous joints to the flagellum, lateral lobes are absent from the head, and the tail appendages are completely to be seen. All the segments of the abdomen are distinctly narrower than those of the thorax and in this it agrees with *Trichoniscus*, *Trichoniscoides*, *Philoscia*, and *Metoponorthus*. In these, however, the flagellum has never more than seven joints, the tail appendages (as in all genera but *Ligia* and *Ligidium*) are partially hidden by the last segment, and in all the four but *Philoscia* there are lobes to the head.

Ligidium hypnorum, Cuvier. [PLATE II.](#)

1792 *Oniscus hypnorum* Cuvier (9), pl. XXVI., figs. 3-5.

1793 *Oniscus agilis* Persoon, quoted by Koch in Panzer (51), part 5, pl. XXIV.

1830 *Ligia hypnorum* Bosc (2), p. 179.

1833 *Ligidium persoonii* J. F. Brandt (3), p. 174, pl. IV., figs. 6-7.

1840 *Zia agilis* Koch (34), part 34, pls. XXII. and XXIII.

1844 *Ligidium personii* Zaddach (77), p. 17.

1853 *Ligidium personii* Lereboullet (39), p. 14, pl. I., fig. 1, pl. II., figs. 20-31.

1857 *Ligidium personii* Kinahan (32), p. 275, pl. XXI., fig. 14, pl. XXII., fig. 9.

1873 *Zia saundersii* Stebbing (70), p. 286.

1873 *Ligidium agile* Norman (48), p. 419.

1885 *Ligidium hypnorum* Budde-Lund (8), p. 254.

1898 *Ligidium hypnorum* G. O. Sars (59), p. 158, pl. LXXI.




FIG. 36.—FLAGELLUM AND LAST PEDUNCULAR JOINT OF THE ANTENNA OF *Ligidium hypnorum*.

This species, which like the last, is the only British representative of its genus, was added to our fauna in 1873 by the Rev. Thomas R. R. Stebbing (70) who found specimens in the neighbourhood of Copthorne Common, Surrey. Up to the present time, when we are pleased to announce that we discovered it in the spring of 1902 at Warley in Essex, *Ligidium hypnorum* has not been recorded from any other place in the British Islands.

As the name of the species implies, it lives in wet situations and in its turn connects *Ligia* with the forms which inhabit drier places. *Ligidium hypnorum* might be mistaken for *Philoscia muscorum*, but as already pointed out in the generic description, the latter has but a few (three) joints to the flagellum, instead of from ten to thirteen. From *Ligia*, the species under consideration is distinguished by its small size, narrow abdomen, and habitat.

BRITISH LOCALITIES:—

England: Warley, Essex; (W.M.W.): Copthorne Common, Surrey; (Stebbing, 70).

FOREIGN DISTRIBUTION:—

Europe: France; (25); Sweden; Denmark; Germany; (59); Turkey; (8).

[Pg 22]

Flagellum with less than ten joints; head with lateral lobes; tail appendages partly hidden.

Genus—**TRICHONISCUS** Brandt, 1833 (3), p. 174.

Abdomen narrow; eyes compound; flagellum, usually with more than three joints.

In *Trichoniscus* the flagellum may have from seven to four (rarely three) joints. As in *Trichoniscoides* and *Haplophthalmus* there are lateral lobes to the head, though these are not very pronounced; the body is also of small size, the abdomen narrow with both divisions of the tail appendages equally so, and almost of the same length though slightly covered by the last segment. The compound eyes distinguish *Trichoniscus* from the two genera named, and from *Platyarthrus*, while its small size and the character of its tail-parts mark it out from all others.

Trichoniscus pusillus Brandt. [PLATE III.](#)

1833 *Trichoniscus pusillus* Brandt (3), p. 174, pl.

IV., fig. 9.

1838 *Itea riparia* Koch (34), part 22, pl. XVII.

1844 *Itea lævis* Zaddach (77), p. 16.

1857 *Philougria celer* Kinahan (32), p. 281, pl.

XXII., figs. 1-4.

1858 *Philougria riparia* Kinahan (33), pp. 191 and

198, pl. XXIII., fig. 1.

1868 *Philougria riparia* Bate and Westwood (1), p.

456.

1898 *Trichoniscus pusillus* Sars (59), p. 161, pl.

LXXII., fig. 1.

This tiny species is found commonly amongst the roots of the herbage in very moist places. It presents a horny translucent appearance and is of a reddish brown colour. It runs with considerable speed, and when it is moving, the white irregular lines with which it is beset are not evident. *Trichoniscus pusillus* is very much like *Trichoniscus vividus* in colour but the latter species is nearly twice as big and has from five to seven joints to the flagellum, while the former has never more than four. *Trichoniscus roseus* is also much larger and its bright red colour (which it loses, however, when preserved in alcohol) is another means of distinguishing it from the species under consideration.

Professor Sars in his *Crustacea of Norway* (p. 162) describes from Christiania, under the name of *Trichoniscus pygmæus*, a still smaller species. As this may possibly be discovered in this country a brief comparison between it and *Trichoniscus pusillus* may be of value. The former reaches a length of but two millimetres; it is "whitish, semi-pellucid with a few light brown pigmentary ramifications across the segments and a double row of irregular opaque patches along the middle of its back" (p. 163). Its body is covered with minute tubercles and there are only three joints to the flagellum; its movements are by no means rapid.

The body of *Trichoniscus pusillus* is smooth and polished. It has four joints to the flagellum—Dr. Scharff (63) says three or four—and it moves quickly.

BRITISH LOCALITIES:—

England: Brightlingsea; Warley; (W.M.W.): Epping Forest; (Bate and Westwood, 1): Hanwell; Southall; Kew Gardens; Langley; Burnham Beeches; Dropmore; Skirmett; Bluebell Hill, Maidstone; (W.M.W.): Chislehurst; Plymouth; Polperro; Looe; (Bate and Westwood, 1): Hertfordshire;




FIG. 37.—FLAGELLUM AND LAST PEDUNCULAR JOINT OF THE ANTENNA OF *Trichoniscus pusillus*.

Northumberland; Durham; (Norman, 49): Exeter; (Parfitt, 53).

Scotland: Edinburgh; (Scott, 68): Cumbrae; (Robertson, 57).

Ireland: Connemara; (Norman, 49): Dublin; Wexford; Cork and Kerry; (Percival Wright *teste* Bate and Westwood, 1): Tyrone; Waterford; Portlaw; Kilkenny; Wicklow; (Kinahan, 33).

FOREIGN DISTRIBUTION:—

Europe: France; (25): Spain; (15): Italy; (19): Norway; Sweden; Denmark; Germany; (59).

Africa: Algeria; Tunis; Azores; (24).

America: Niagara; North America; (59).

Trichoniscus vividus, Koch. [PLATE IV.](#) (from a spirit specimen).

1840 *Itea vivida* Koch (34), part 34, pl. IV.

1858 *Philougria vivida* Kinahan (33), pp. 197 and 198, pl. XXIII., fig. 2.

1868 *Philougria vivida* Bate and Westwood (1), Vol. II., pp. 458 and 459, figs.

This species is claret-brown in colour and under a lens it is seen to be marbled with white, indeed in appearance it is much like *Trichoniscus pusillus* though twice the size. There are important differences between the two species as regards the number of joints to the flagellum. These vary from five to seven in *Trichoniscus vividus* while in the other, as already pointed out, there are not more than four.

[Pg 24]




FIG. 38.—FLAGELLUM
AND LAST PEDUNCULAR
JOINT OF THE ANTENNA
OF *Trichoniscus*
vividus.

The body is practically speaking smooth for it bears only very small tubercles, widely separated. In *Trichoniscus vividus* the antennæ lack the bristles which characterise those of the other species in the genus. The species under consideration was discovered by Dr. Kinahan in March, 1858, at Portlaw, Co. Waterford and is active even amongst the snow.

BRITISH LOCALITIES:—

Ireland: Portlaw, Co. Waterford; (Kinahan, 33): Cappagh, Co. Waterford; (Scharff, Irish Nat., Vol. IX., p. 158): Borris, Co. Carlow; (Scharff, 64.)

FOREIGN DISTRIBUTION:—

Europe: Spain; (12).

Trichoniscus roseus Koch. [PLATE V.](#)

1838 *Itea rosea* Koch (34), part 122, pl. XVI.

1858 *Philougria rosea* Kinahan (33), pp. 197 and 199, pl. XXIII., fig. 3.

1858 *Philougria rosea* Bate and Westwood (1), p. 460.

1898 *Trichoniscus roseus* Sars (59), p. 163, pl. LXXIII., fig. 1.

The third British species of *Trichoniscus* is of a deep pink colour and has a light yellow stripe down the back (in some habitats the animals are said to be quite white). Arranged in transverse rows upon the body are large tubercles, each of which under strong magnification will be found to end in a tiny hair. It is distinguished from *Trichoniscus pusillus* by the larger size of its body, which is also comparatively broader, and from *Trichoniscus vividus* by the four joints of the flagellum of its antennæ which latter have strong bristles upon them. In the former species there are five or more joints to the flagellum and the antennæ, though hairy, lack the bristles. *Trichoniscus roseus* is to be looked for in old gardens.

BRITISH LOCALITIES:—

England: Warley; (W.M.W.): Maldon; (W.M.W. from R.M.): Stanmore; Hanwell; Ealing; Wimbledon; (W.M.W.): Berkhamsted; Torquay; (Norman, 49): Plymouth; (Bate and Westwood, 1 and B.M.): Grassendale, near Liverpool; (R.W.): Newtownards; (R.W., Irish Nat, 1904, p. 260.)

Scotland: Tarbert; (Scot, 68).

Ireland: Dublin; Ballyfinder, Co. Down; (Scharff, 63): Templeogue; Dundrum; Blackrock; Rathgar, Co. Dublin; Bray, Co. Wicklow; (R.F.S.): Oakleigh; Kerry; (R.W.): Belfast; (Welch, Irish Nat., 1896, p. 213.): At the grave of Josiah Welch (grandson of John Knox), Castle Upton; Richhill, Co. Armagh; Castleconnell Ferry; (R.W.): Glenade House, Co. Antrim; (R.W. from R. Ll. Praeger).

FOREIGN DISTRIBUTION:—

Europe: France; (25): Spain; (12): Italy; (59): Denmark; Germany; Holland; (39): Dalmatia; (18).

Africa: Algeria; Tunis; (24).

Genus—**TRICHONISCOIDES**, Sars, 1898 (59), p. 164.

Abdomen narrow; eyes simple; (or wanting); flagellum, with four joints.

The members of this genus are very much like those of *Trichoniscus*. In the latter, however, the hinder legs are longer in proportion and the eyes are compound.

Trichoniscoides albidus Budde-Lund.

PLATE VI.

1879 *Trichoniscus albidus* Budde-Lund (7) p. 9.

1898 *Trichoniscoides albidus* Sars (59), p. 165, pl. LXXIII., fig. 2.

We are able to include this species, as a specimen was found by Mr. Webb at Eton Wick in the summer of 1899. It is one of a number of species which the Rev. Canon Norman (49, p. 18) suggested as likely to be British. It is the only representative of its genus, which does not differ in any very important characters from the others in the family. The narrow elongated body will serve to separate it from *Trichoniscus vividus* and *Trichoniscus roseus*, but on account of its size, which is much the same as that of *Trichoniscus pusillus* and the two British species of *Haplophthalmus*, it will be advisable to give some further points of distinction. From the first its white colour will serve to differentiate it; the other two lack the narrow abdomen seen in *Trichoniscoides albidus*. Moreover, not one of the three shows the serrations on the side plates which characterise the species under consideration. *Platyarthrus hoffmannseggii* is small and white and the edges of its side plates are toothed, but it is oval in shape, possesses no eyes, and its stout antennæ have but a single joint to the flagellum instead of four. On the Continent this species has been found in rich soil.

BRITISH LOCALITIES:—

England: Eton; (Stebbing, 71a): Sunderland; (Brady, 50a).

FOREIGN DISTRIBUTION:—

Europe: France; Wimereux and Lyons, Forêt (25): Norway; Denmark; (59).




FIG. 39.—FLAGELLUM AND LAST PEDUNCULAR JOINT OF THE ANTENNA OF *Trichoniscus roseus*.




FIG. 40.—FLAGELLUM AND LAST PEDUNCULAR JOINT OF THE ANTENNA OF *Trichoniscoides albidus*.

Genus—**HAPLOPHTHALMUS** Schöbl, 1850 (66), p. 449.

Abdomen broad (comparatively); eyes simple; flagellum with three joints; back with longitudinal ridges.

The body of *Haplophthalmus* is long in proportion to its width, but there is no abrupt decrease in the breadth of the abdomen as seen in *Trichoniscus* and *Trichoniscoides*. The eyes are simple as in the latter genus and the lateral lobes of the head are rather large, while the side plates of the body are well separated.

Haplophthalmus mengii Zaddach. [PLATE VII.](#)

1844 *Itea mengii* Zaddach (77), p. 16.

1860 *Haplophthalmus elegans* Schöbl (66), p. 449.

1885 *Haplophthalmus mengii* Budde-Lund (8), p. 250.

1898 *Haplophthalmus mengii* Sars (59), p. 167, pl. LXXIV., fig. 1.




FIG. 41.—FLAGELLUM
AND LAST PEDUNCULAR
JOINT OF THE ANTENNA
OF *Haplophthalmus
mengii*.

The Rev. Canon Norman discovered two specimens of this species in Ireland in June, 1900 (50); in the previous year one of us (Mr. Webb) found a single example at Eton Wick.

The main differences between the members of this genus and their allies are set forth in the generic description and incidentally elsewhere, so we shall content ourselves with giving the distinctive points of the two British species. *Haplophthalmus mengii* has a number of raised longitudinal ribs on each segment of the thorax, the outer ridges being somewhat broken. There are also two prominent ribs upon the third segment of the abdomen.

BRITISH LOCALITIES:—

England: Eton; (Stebbing, 71a);
Sunderland; (Brady, 50a).

Ireland: Corcumroe Abbey; Co. Clare (Norman, 50).

FOREIGN DISTRIBUTION:—

Europe: France; (25); Norway; Prussia; Germany; Bohemia; (59).

Haplophthalmus danicus Budde-Lund.
[PLATE VIII.](#)

1870 *Haplophthalmus elegans* Budde-Lund (6), p. 228 (not *Haplophthalmus elegans* Schöbl).

1879 *Haplophthalmus danicus* Budde-Lund (7), p. 9.

1881 *Haplophthalmus mengii* Weber (76), p. 192, pl. V., figs. 7-9 (not *Itea mengii* Zaddach).

1885 *Haplophthalmus danicus* Budde-Lund (8), p. 250.

1898 *Haplophthalmus danicus* Sars (59), p. 168, pl. LXXIV., fig. 2.




FIG. 42.—FLAGELLUM
AND LAST PEDUNCULAR
JOINT OF THE ANTENNA
OF *Haplophthalmus
danicus*.

This species was added to the British list by the Rev. Canon Norman (49), who found a colony in his garden at Berkhamsted. It has rows of tubercles on its thorax instead of ridges, and there are no ribs at all upon the abdomen. The front of the head projects further comparatively and forms a more acute point than in *Haplophthalmus mengii* and it is not so purely white in colour as the latter species.

BRITISH LOCALITIES:—

England: Warley Place; (W.M.W. from Miss Willmott); Queen's Cottage, Kew Gardens; Stanmore; Hanwell, garden

at Odstock, Bennett's Nurseries; (W.M.W.): Berkhamsted; (Norman, 49): Sunderland; (Brady, 50a).

FOREIGN DISTRIBUTION:—

Europe: France; (25): Denmark; Holland; Germany; (Dollfus, Feu de Jeun, Nat., April, 1896): Norway; (Sars, 59).

Section—ONISCI.

THE OUTER DIVISIONS OF THE TAIL APPENDAGES
BROADER THAN THE INNER ONES.

Family—ONISCIDÆ.

Tail appendages projecting when the animal is walking.

(1.) Unable to roll up into a complete ball.

Genus—ONISCUS Linné 1746 (41), p. 360.

Flagellum, with three joints; abdomen broad; head with lateral lobes.

The characters given above taken in conjunction with the size of the animals will serve to distinguish the members of this genus.

Oniscus asellus Linné (The "Common Slater.") [PLATE IX.](#)

1761 *Oniscus asellus* Linné (41), p. 500, No. 2058.

1792 *Oniscus murarius* Cuvier (9), p. 22, pl. XXVI.

1838 *Oniscus fossor* Koch (34), part 22, pl. XXII.

1868 *Oniscus asellus* Bate and Westwood (1), p. 468.

1868 *Oniscus fossor* Bate and Westwood (1), pp. 471-2.

1898 *Oniscus asellus* Sars (59), p. 171, pl. LXXV.

Oniscus asellus is one of the largest of our woodlice and it is also probably the commonest, though *Porcellio scaber* is in many places quite as abundant. The body of *Oniscus* is broad and expanded and the colour is usually a slate grey with yellowish markings more or less regularly arranged.

[Pg 28]




FIG. 43.—FLAGELLUM
AND LAST PEDUNCULAR
JOINT OF THE ANTENNA
OF *Oniscus asellus*.

From the genus *Porcellio* the species with which we are concerned is at once distinguished by the three jointed flagellum. *Porcellio* has but two joints and has, besides, a prominent lobe projecting from the middle of the head, which is not seen in *Oniscus*. *Philoscia*, although it has three joints to the flagellum, has a narrow abdomen and lacks entirely the lateral lobes which are a feature of the other genera of Oniscidæ.

Oniscus fossor of Koch (34) was recognized by Kinahan and by Bate and Westwood as a species. Dr. Scharff submitted specimens to Professor Budde-Lund who found no differences between them and *Oniscus asellus*. The former (63) mentions, however, that the characteristics of the supposed species are those of young examples of *Oniscus asellus*, and Professor Sars (59, p. 173) seems to be of the same opinion. Many young examples of *Oniscus asellus* that we have examined have a curious whitish transverse band owing to the light colour of the dorsal plates of the first abdominal segments. The flagellum also does not seem to shew in young animals a distinct division into three joints.

BRITISH LOCALITIES:—

England: High Beach, Epping, including an albino; Maldon; Brightlingsea; Iver; Hanwell; Eton; Kew; Pamber Forest; Kingston-on-Soar; Bluebell Hill, Maidstone;

(W.M.W.): Lynmouth; (W.M.W. from J.T.C.).

Scotland: (Scharff, 63). Dinnet, Aberdeenshire; (W.M.W. from Madame Christen).

Ireland: (Scharff, 63). Yellow form with black spots, Donegal (R.W.)

FOREIGN DISTRIBUTION:—

Europe: Almost throughout; (12): France; (25): Spain; (12): Sweden Norway; Denmark; Germany; Holland; Italy; Iceland; (59): Faroe Islands; Thorsharn; (R.F.S.)

Africa: Azores; (24).

America: Greenland; (59): North America; (Budde-Lund).

Genus—**PHILOSCIA** Latreille, 1804 (37), p. 43.

Flagellum with three joints; abdomen narrow; head without lateral lobes.

If any further differences of an obvious kind be required to distinguish *Philoscia* from *Oniscus*, one at least will be found in the much greater development of the hinder legs in the former genus.

Philoscia muscorum Scopoli. [PLATE X.](#)

[Not of Lereboullet, which is an *Oniscus*, see Bate and Westwood (1).]

1763 *Oniscus muscorum* Scopoli (67), p. 415.

1793 *Oniscus sylvestris* Fabricius (26), p. 397.

1793 *Oniscus agilis* Koch in Panzer (51), part 9, pl. XXIV.

1833 *Philoscia marmorata* Brandt (3), p. 183.

1838 *Ligia melanocephala* Koch (34), part 22, pl. XVIII.

1847 *Zia melanocephala* Koch (38), part 40, pl. I. p. 212.

1868 *Philoscia muscorum* Bate and Westwood (1), p. 450.

1898 *Philoscia muscorum* Sars (59), p. 174, pl. LXXVI., fig. 1.

This species lives chiefly at the roots of grass and under the stones or sticks that lie among it. *Philoscia muscorum* has a very smooth and shining body, and its long legs enable it to move very rapidly. The ground colour of its dorsal surface varies from light yellow to deep brown. There are characteristic dark markings down the middle of the thorax and on the sides, between which are lighter patches. In dark coloured specimens the markings are by no means so evident.

BRITISH LOCALITIES:—

England: High Beach, Epping; Warley; (W.M.W.): Maldon; (W.M.W. from R.M.): Kew; Langley; Hanwell, yellow variation; Bluebell Hill, Maidstone; (W.M.W.): Liphook; (C.S.): Pamber Forest; Kingston-on-Soar; (W.M.W.)

Scotland: (Scott, 68).

Ireland: Almost throughout; (Scharff, 63).

FOREIGN DISTRIBUTION:—

Europe: France; (25): Spain; (12): Sicily; (19): Hertsogovina; (22): Sweden; (21): Norway; Denmark; Prussia; Germany; Holland; Poland; Austria; Italy; (59): Sardinia; (21).

Africa: Algeria; Tunis; (24).

Philoscia couchii Kinahan. [PLATE XI.](#)

1858 *Philoscia couchii* Kinahan (33), p. 195, pl. XXIII., fig. 4.




FIG. 44.—FLAGELLUM AND LAST PEDUNCULAR JOINT OF THE ANTENNA OF *Philoscia muscorum*.

- 1868 *Philoscia couchii* Bate and Westwood (1), p.
 1885 *Ligidium couchii* Budde-Lund (8), p. 257.
 1885 *Philoscia longicornis* Budde-Lund (8), p. 221.
 1897 *Philoscia couchii* Dollfus (21), p. 72, pl. I., fig.
 1.




FIG. 45.—FLAGELLUM
 AND LAST PEDUNCULAR
 JOINT OF THE ANTENNA
 OF *Philoscia couchii*.

Philoscia couchii is an inhabitant of the sea-side; it is smaller than the last species, its colour to the naked eye is a uniform lead-grey, and its antennæ are very large (compared with its size) and hairy.

This species was discovered by Professor Kinahan when in the company of Messrs. Bate and Westwood near Polperro in Cornwall in the year 1858, and dried specimens presented by him are in the British Museum (Natural History).

BRITISH LOCALITIES:—

England: Talland Cove; Polperro; (Bate and Westwood, 1): Salcombe, Devon; (Norman, 49): Meadefoot, Torquay; (Stebbing in 49).

FOREIGN DISTRIBUTION:—

Europe: France; (25): Spain; (12): Sicily; (19): Sebastopol; (Norman, 49).

Africa: Azores; Canaries; Morocco; Algiers; Tunis: Egypt Senegal; (24).

Atlantic Isles: Canaries; Azores; (21).

Asia: Syracuse; Bazone (18).

Genus—PLATYARTHURUS Brandt, 1833 (3), p. 174.

[*Typhloniscus* Schöbl (66), p. 279.]

Flagellum with one joint; eyes wanting; abdomen broad; habitat, ants' nests.

The broad body, which is much flattened, and the very thick antennæ distinguish *Platyarthrus* from the other small woodlice (Trichoniscidæ).

Platyarthrus hoffmannseggii Brandt. [PLATE XII.](#)

1833 *Platyarthrus hoffmannseggii* Brandt (3), p. 174, pl. IV., fig. 10.

1844 *Itea crassicornis* Koch (34), part 36, pl. V.

1860 *Typhloniscus steinii* Schöbl (66), p. 282.

1868 *Platyarthrus hoffmannseggii* Bate and Westwood (1), p. 464.

1898 *Platyarthrus hoffmannseggii* Sars (59), p. 175, pl. LXXVI., fig. 2.

Up to the present this is the only woodlouse which has been found in the nests of British ants. It is small and oval, its colour is white, and its body is covered with tubercles. The edges of its side plates are toothed, its flagellum has but a single joint and it has no eyes.

Miss Kate Hall tells us that, if very hungry, ants in captivity will kill and eat *Platyarthrus*. With regard to its own food, Lord Avebury has favoured us with the opinion that it lives on the spores of the lower plants, such as would be found in the ants' nest.

BRITISH LOCALITIES:—

England: Warley; Hanwell; West Drayton; Langley; Kingston-on-Soar; Bluebell Hill, Maidstone; (W.M.W.): Berkhamsted; Salcombe; Devon; Cheddar Cliffs, Somerset; (Norman, 49): Ide, near Exeter; (Parfitt, 53): Torquay; (Stebbing in 49); Lulworth Cove; (Rev. A. R. Hogan *teste* Bate and Westwood, 1): Hammersmith; Oxford; Berry Head, Torquay; Plymouth; (Bate and Westwood, 1): In the nest of *Myrmica rubra*, Newton Ferrers (E. E. Lowe).




FIG. 46.—FLAGELLUM
AND LAST PEDUNCULAR
JOINT OF THE ANTENNA
OF *Platyarthrus*
hoffmannseggii.

Scotland: Banff; (Thomas Edward in 49).

Ireland: Leixlip, Co. Dublin; Lismore, Co. Waterford; Glengariff, Co. Cork; (Scharff, 63): Bagenalstown, Co. Carlow; (64).

FOREIGN DISTRIBUTION:—

Europe: France; (28); Spain; (12); Denmark; Germany; Holland; Bohemia; Austria; Tyrol; Helvetia; (59).

NOTE.—In the genera which follow, air-tubes or air-cavities (tracheæ) are present in the outer plates of the abdominal appendages, 1 and 2, or 1 to 5. The appendages in question have in consequence a milk-white appearance in the living animal owing to the fact that the enclosed air reflects white light. Considerable interest attaches to the study of these tracheæ, which have the same function as those of insects, but which have been independently developed. To emphasise the latter fact the structures are often termed "pseudotracheæ."

[Pg 32]


Genus—**PORCELLIO** Latreille, 1804 (37), p. 45.

Flagellum, with two joints; *abdomen*, broad; *frontal lobe* projecting.

Porcellio is easily separated from the previous genera—*Oniscus*, *Philoscia*, and *Platyarthrus*—by its two-jointed flagellum. The fact that the abdomen is not abruptly narrowed separates it from *Metoponorthus*, which also lacks the prominent frontal lobe so characteristic of *Porcellio*. The species of this genera might be confused with *Cyclisticus* which has two joints to the flagellum and a broad abdomen, but the latter genus has the power of rolling itself into a ball, while its frontal lobe is very small, and the first segment of its thorax is comparatively larger than in any species of *Porcellio*.

Porcellio scaber Latreille. [PLATE XIII.](#)

- 1804 *Porcellio scaber* Latreille (37), p. 45.
- 1818 *Oniscus granulatus* Lamarck (36), p. 261.
- 1818 *Porcellio nigra* Say (62), p. 432.
- 1840 *Porcellio brandtii* Milne-Edwards (46), p. 168.
- 1840 *Porcellio dubius* Koch (34), part 34, pl. VIII.
- 1847 *Porcellio asper* Koch (35), p. 207, pl. VIII., fig. 98.
- 1857 *Porcellio montezumæ* Saussure (60), p. 207.
- 1865 *Porcellio paulensis* Heller (31), p. 136, p. XII., fig. 5.
- 1868 *Porcellio scaber* Bate and Westwood (1), p. 475.
- 1876 *Porcellio graniger* Miers (44), p. 223.
- 1885 *Porcellio graniger* Budde-Lund (8), p. 149.
- 1898 *Porcellio scaber* Sars (59), p. 176, pl. LXXVII.



The body of *Porcellio scaber* is densely covered with tubercles. Its colour is usually of a very dark grey, but at times it is quite red or variegated with yellow. Albino specimens have been recorded. The two joints of the flagellum are of the same length and together equal that of the last joint of the peduncle. Air-tubes are present in the outer plates of the first two abdominal appendages.

BRITISH LOCALITIES:—

England: High Beach, Epping; Warley; Brightlingsea; (W.M.W.): Maldon; (W.M.W. from R.M.): Langley; Kew; Skirmett; Pamber Forest; (W.M.W.):

FIG. 47.—FLAGELLUM AND LAST PEDUNCULAR JOINT OF THE ANTENNA OF *Porcellio scaber*. Liphook; (C.S.): Stoke-on-Trent; Kingston-on-Soar; (W.M.W.) Scotland: Dinnet (W.M.W. from Madame Christen).

Ireland: Common everywhere; (Scharff, 63.)

FOREIGN DISTRIBUTION:—

Europe: Throughout; (59): France; (28): Spain; (15): Iceland; (59): Faroe Isles—Thorsharn and Naalsole—(R.F.S. and B.M., N. Annadale).

America: Greenland; North America; Sandwich Isles; (B.M.); Mexico; (59): St. Paul; St. Croix; (59); Ascension; Tristan d'Acunha; (23)

Asia: Ceylon; Kamtschatka: (23).

Australia: Melbourne; Sydney; Tasmania; New Zealand; (B.M., Chilton).

Africa: Azores; Canaries; Cape of Good Hope; (24).

Porcellio pictus Brandt and Ratzeburg. [PLATE XIV.](#)

1833 *Porcellio pictus* Brandt and Razteburg (4), p. 78, pl. 12, fig. 5.

1839 *Porcellio melanocephalus* Koch (34), part 28, pl. XVIII.

1853 *Porcellio melanocephalus* Schnitzler (65), p. 24.

1856 *Porcellio mixtus* Fitch (29), p. 120.

1868 *Porcellio pictus* Bate and Westwood (1), p.

1898 *Porcellio pictus* Sars (59), p. 177. pl. LXXVII., fig. 1.

There are tubercles on the body of *Porcellio pictus*, which is a striking looking animal. Its head is black with the lateral lobes curved outwards; there is a dark band down the middle of the back and commonly two others on each side, with more or less conspicuous yellow markings between.

The distal (terminal) joint of the flagellum is but half the length of the proximal one and the last peduncular joint is longer than the two combined.

The abdominal appendages—1 and 2—are provided with air-tubes.

BRITISH LOCALITIES:—

England: Maldon; (W.M.W. from R.M.): Chislehurst; (Bate and Westwood, 1): Cooper's Hill, near Cheltenham; (Norman, 49): Exeter; (Parfitt, 53): Kent; (Bate and Westwood, 1.)

Scotland: Between Leith and Portobello; (Scott, 68): Cumbrae (Scott, 68a): Ayrshire; (Boyd in Norman, 49): Banff; (T. Edwards in Norman, 49).

Ireland: Dublin; Belfast; (Bate and Westwood, 1): Galway; Maryborough; Queen's Co., Castel; and Caher Co. Tipperary; (R.F.S.)

FOREIGN DISTRIBUTION:—

Europe: France; (25): North, West-Central, and East Europe; (8): Sweden; Norway; Denmark; Germany; Hungary; Russia; (59).

North America: (8).

Porcellio dilatatus Brandt. [PLATE XV.](#)

1833 *Porcellio dilatatus* Brandt and Ratzeburg (4), p. 78, pl. 12., fig. 6.

1840 *Porcellio scaber* Milne-Edwards (not Latreille) (46), p. 167.

1868 *Porcellio dilatatus* Bate and Westwood (1), p.

1898 *Porcellio dilatatus* Sars (59), p. 179, pl. LXXVII., fig. 2.




FIG. 48.—FLAGELLUM AND LAST PEDUNCULAR JOINT OF THE ANTENNA OF *Porcellio pictus*.




FIG. 49.—FLAGELLUM
AND LAST PEDUNCULAR
JOINT OF THE ANTENNA
OF *Porcellio dilatatus*.

The fact that *Porcellio dilatatus* is more than half as broad as it is long, at once distinguishes it from the other species of *Porcellio*. It is tuberculated and of somewhat a lighter grey than *Porcellio scaber* usually is. The two species agree in having the two joints of the flagellum equal, but the last peduncular joint, as in *Porcellio pictus*, is longer than the flagellum.

As in the two preceding species, air-tubes are found in the outer plates of the appendages on the first two abdominal segments. *Porcellio dilatatus* is to be looked for near houses.

BRITISH LOCALITIES:—

England: Maldon; (W.M.W. from R.M.): Eton; (Stebbing from W.M.W., 71a): Berkhamsted; (Norman, 50): Headley, Surrey; Ventnor; (Stebbing in Norman,

49).

Ireland: Dublin; (Scharff, 63): Dundrum; (Scharff in Norman, 50): Galway; Roundstone; (R.F.S.): Belfast; (C. W. Buckle, *Irish Nat.*, Vol. XI. (1902), p. 43).

FOREIGN DISTRIBUTION:—

Europe: France; (25): Spain; (12): Denmark; Norway; Germany; Poland; Holland; (59).

Africa: Madeira; Azores; (24).

Australia: New Guinea; (59).

***Porcellio rathkei* Brandt. [PLATE XVI.](#)**

1833 *Porcellio rathkei* Brandt (3), p. 177, fig. 10.

1833 *Porcellio ferrugineus* Brandt (3), p. 178.

1840 *Porcellio trilineatus* Koch (34), part 34, pl. IX.

1853 *Porcellio trivittatus* Lereboullet (39), p. 54, pl. I., figs. 13 and 14.

1853 *Porcellio tetramoerus* Schnitzler (65), p. 24.

1853 *Porcellio striatus* Schnitzler (65), p. 24.

There is often a light band down the back and one on either side of it near the margin in *Porcellio rathkei* (especially in the males), with other more irregularly arranged light patches between. Unlike the three species previously considered, the present one has a smooth body. The distal joint of the flagellum is the longer, and the flagellum itself is equal in length to the last joint of the peduncle.

Some specimens found by Mr. Webb in 1899 at Eton were submitted to Mr. Stebbing, and since then the former has found *Porcellio rathkei* to be pretty generally distributed in West Middlesex, where the species appears to frequent the open fields.

Air-tubes occur in abdominal appendages 1 to 5 and the white appearance of all of these at once serves to distinguish the living animal from *Porcellio scaber* in which the first two pairs of abdominal appendages alone are white.

BRITISH LOCALITIES:—

England: Eton; (Stebbing, 71a): Lane End; (Stebbing, from the Misses Johnston, 71a): Acton; Ealing; Hanwell; Southall; Northolt; Greenford; West Drayton; Mortlake; (W.M.W.); Sunderland; (Brady, 50a).

FOREIGN DISTRIBUTION:—

Europe: France; (25): Bosnia; Servia; (22): Hertzogovania (B.M.); Norway;




FIG. 50.—FLAGELLUM
AND LAST PEDUNCULAR
JOINT OF THE ANTENNA

Asia: Transcaucasia; (59).

North America: (59).

Porcellio laevis Latreille. [PLATE XVII.](#)




FIG. 51.—FLAGELLUM
AND LAST PEDUNCULAR
JOINT OF THE ANTENNA
OF *Porcellio laevis*.

1804 *Porcellio laevis* Latreille (37),
p. 46.

1827 *Porcellio degeerii* Savigny
and Audouin (61), p. 289.

1833 *Porcellio cucercus* Brandt (3),
p. 177.

1833 *Porcellio syriacus* Brandt (3),
p. 178.

1833 *Porcellio musculus* Brandt
(3), p. 180.

1833 *Porcellio cinerascens* Brandt
(3), p. 178.

1833 *Porcellio dubius* Brandt (3),
p. 178.

1837 *Porcellio poeyi* Guérin (30), p.
6.

1844 *Porcellio urbicus* Koch (34),
part 36, pl. IV.

1847 *Porcellio flavipes* Koch (35),
p. 206, pl. VIII., fig. 97.

1853 *Cylisticus laevis* Schnitzler
(65), p. 25.

1857 *Porcellio cubensis* Saussure (60), p. 307.

1857 *Porcellio sumichtasli* Saussure (60), p. 307.

1857 *Porcellio cotillæ* Saussure (60), p. 307.

1857 *Porcellio aztecus* Saussure (60), p. 307.

1857 *Porcellio mexicanus* Saussure (60), p. 307.

Another smooth species is *Porcellio laevis*. The colour of its body is light grey with irregular white markings. The large size of this species and the very long tail-appendages of the males are features which will help to identify it. The distal joint of the flagellum is slightly the longer and as in the last species (*P. rathkei*) the flagellum is equal in length to the last joint of the peduncle. The chief habitats for this species are among vegetable rubbish near human dwellings.

Only the first two abdominal appendages contain air-tubes.

BRITISH LOCALITIES:—

England: Maldon; (W.M.W. from R.M.): Ipswich (1892); Hanwell; Wimbledon; (W.M.W.): Kent; (Kinahan, 32).

Ireland: Dublin; (Bate and Westwood, 1): Blackrock Dundrum; Co. Dublin; Galway; (R.F.S.)

FOREIGN DISTRIBUTION:—

Europe: France; (25): Spain; (15): Sicily; (19): Hertzogovania; (22): Sweden; Denmark; Germany; Belgium; Austria; Italy; Dalmatia; Greece; Turkey; (59); Corfu; (B.M.): Inca, Majorca (23); (B.M.—Pocock and Thomas.)

Asia: Syria; Turkestan; (21).

Africa: Morocco; Algeria; Tunis; Tripoli; Senegal; Egypt; (23).

Atlantic Isles: Bermudas; Azores; Canaries; Cape Vera; Madeira; (24).

America: North America; Mexico; Peru; Brazil; Chili; West Indies; Pacific Islands; (59); Sandwich Isles; (B.M.)

Porcellio ratzeburgii Brandt. [PLATE XVIII.](#)

1833 *Porcellio ratzeburgii* Brandt, (3), p. 178.

1839 *Porcellio nemorensis* Koch (34), part 28, pl.
XIX.

1839 *Porcellio lugubris* Koch (34), part 28, pl. XX.

1853 *Porcellio quercum* Schnitzler (65), p. 24.

1898 *Porcellio ratzeburgii* Sars (59), p. 182, pl.

LXXX, fig. 1.

There are granulations on the middle of




FIG. 52.—FLAGELLUM
AND LAST PEDUNCULAR
JOINT OF THE ANTENNA
OF *Porcellio ratzeburgii*.

the segments in *Porcellio ratzeburgii* and the sides of its body are more nearly parallel than in the other species of *Porcellio*; the frontal lobe is, practically speaking, semicircular and the lateral plates of the thorax flank the head to a considerable extent. As in *Porcellio pictus*, the dark band is in the middle of the back. The distal joint of the flagellum is nearly twice as long as the proximal, and the flagellum is shorter than the last joint of the peduncle. This species was added to the British list by Mr. Webb (74) in 1898.

Porcellio ratzeburgii agrees with *Porcellio rathkei* in having air-tubes in the first five abdominal appendages.

BRITISH LOCALITIES:—

England: Warley; Brightlingsea; young examples (W.M.W.): Maldon; young examples (W.M.W. from R.M.)

FOREIGN DISTRIBUTION:—

Europe: Trafoi St. Martini, and Capitello, in the Tyrol; (Norman, 50); East Alps, very common; Val-de-Joux; Massif de la Chartreuse Vaulnaveys (25); Bosnia; (22); Norway; Central Europe; Upper Pfaltz; Bohemia; Saxony; Rhaetia; (59).

Genus—**METOPONORTHUS** Budde-Lund, 1879 (7), p. 4.
Porcellionides Miers, 1876 (44), p. 98.

Flagellum, with two joints; abdomen, narrow; frontal lobe not developed.

The hinder legs of *Metoponorthus* are proportionately longer than in any other Oniscidæ save *Philoscia*. Both genera have a narrow abdomen, but *Philoscia* has an extra joint to the flagellum, and shows no sign of lateral lobes to the head.

Metoponorthus pruinus Brandt. [PLATE XIX.](#)

- 1833 *Porcellio pruinus* Brandt (3), p. 181.
- 1840 *Porcellio truncatus* Milne-Edwards (46), p. 173.
- 1840 *Porcellio maculicornis* Koch (34), part 34, pl. XVI.
- 1853 *Porcellio frontalis* Lereboullet (39), p. 63, pl. I, fig. 17.
- 1868 *Porcellio pruinus* Bate and Westwood (1), p. 487.
- 1877 *Porcellio (Porcellioides) flavo-vittatus* Miers (45), p. 669 pl., LXVIII., fig. 4.
- 1898 *Metoponorthus pruinus* Sars (59), p. 184, pl. LXXX., fig. 2.

Undamaged specimens of *Metoponorthus pruinus* are of a beautiful bluish-grey colour, owing to a "bloom" which is easily brushed off, revealing a dark reddish-brown tint beneath it. The antennæ are long and have white markings upon them.

Air-tubes occur in the first two abdominal appendages.

BRITISH LOCALITIES:—

England: Maldon; (W.M.W. from R.M.): Hanwell; Eton Wick; Kew; Ipswich; Stoke-on-Trent; (W.M.W.): Chiselhurst; Oxford; (Bate and Westwood): Berkhamsted; Burnmoor; Durham; (Norman, 49): Exeter; (Parfitt, 53); Torquay; (B.M.—T.R.R.S.)




FIG. 53.—FLAGELLUM
AND LAST PEDUNCULAR
JOINT OF THE ANTENNA

Scotland: Banff; (Thomas Edwards in Norman, 49) OF *Metoponorthus*
pruinus.

Ireland: Dublin; (Kinahan, 32): Foyle District; Donegal; Galway; Clonbrock, Co. Galway; Mornington, Co. Meath; Santry; Gleeson Park; Dundrum, Co. Dublin; Bray; (R.F.S.)

FOREIGN DISTRIBUTION:—

Europe: Practically all the Countries of Europe are given in Dollfus' list; (23).

Asia: Japan; China; Syria; Ceylon; Sumatra; Celebes; Phillipines; Caucasus; Himalayas; (23): Christmas Island; (B.M.)

Africa: Generally distributed; Madagascar; Seychelles; (23).

Atlantic Isles; (23).

America: North and South, almost everywhere, to judge from M. Dollfus' list; (23).

Australia: New Caledonia; (23).

Metoponorthus cingendus Kinahan. [PLATE XX.](#)

1857 *Porcellio cingendus* Kinahan (32), p. 279, pl. XIX., figs. 1468-9.

1868 *Porcellio cingendus* Bate and Westwood (1), p. 489.

1885 *Metoponorthus simplex* Budde-Lund (8), p. 188.




FIG. 54.—FLAGELLUM
AND LAST PEDUNCULAR
JOINT OF THE ANTENNA
OF *Metoponorthus*
cingendus.

The colour of *Metoponorthus cingendus* is steel blue with red or yellowish spots. It has a raised line across each thoracic segment and its abdomen is narrower than in *Metoponorthus pruinus*.

BRITISH LOCALITIES:—

England: Salcombe, Devon; (Norman, 49): South Devon; (Stebbing in 49).

Ireland: Dublin; (B.M. from Kinahan); Mountain Districts of Dublin, Wicklow, and Cork; Coast of Kerry; Arran Islands; Achill, Co. Mayo; Roundstone, Co. Galway; Mallow, Caef Island; Glandore; Brock Haven, Co. Cork; Killoughrim Forest, Co. Wexford; Kenmare, Co. Kerry; (R.F.S.).

FOREIGN DISTRIBUTION:—

Europe: France; (25): Spain; (12).

(2.) Able to roll up into a ball.

Genus—CYLISTICUS Schnitzler, 1853 (65), p. 24.

Flagellum, with two joints; *abdomen broad*; *frontal lobe*, very small.

The characters given immediately above are almost those of *Porcellio* with which *Cylisticus* might, perhaps, be confounded. The latter has the power, however, of rolling itself into a ball, and the first segment of the thorax is comparatively larger than in any species of *Porcellio*, indeed the side plates of the segment in question entirely flank the head. These features, as well as the straight sides of the body and the arched back, connect *Cylisticus* with *Armadillidium*, from which the former is, however, at once separated by its long pointed tail appendages.

Cylisticus convexus De Geer. [PLATE XXI.](#)

1778 *Oniscus convexus* De Geer (10), p. 553, pl. XXXV., fig. 11.

1833 *Porcellio spinifrons* Brandt (3), p. 177.

1836 *Porcellio laevis* Koch (34), part 6, pl. I.

1853 *Porcellio armadilloides* Lereboullet (39), p.

65. pl. I., fig. 18.
 1853 *Cylisticus laevis* Schnitzler (65), p. 25.
 1868 *Porcellio armadilloides* Bate and Westwood
 (1), p. 485.
 1898 *Cylisticus convexus* Sars (59), p. 186, pl.
 LXXXI.

There is but a single species of *Cylisticus* found in this country, so that it is not necessary for us to go into much further detail with regard to it. *Cylisticus convexus* has the two joints of the flagellum about equal, and they together in turn closely approximate in length to the last joint of the peduncle. Mr. Stebbing says, in a letter, that British examples do not appear to have the "white tail-piece" seen in Continental ones. It is not noticeable in the preserved specimens which we have seen from Berkhamsted and Leixlip, but it is very evident in the living ones found at Hanwell and Maidstone.




FIG. 55.—FLAGELLUM
 AND LAST PEDUNCULAR
 JOINT OF THE ANTENNA
 OF *Cylisticus convexus*.

The abdominal appendages 1 to 5 are provided with air-tubes.

BRITISH LOCALITIES:—

England: Maldon; (W.M.W. from R.M.): Hanwell; Bluebell Hill, Maidstone; Eton; (W.M.W.): Berkhamsted; Portland; (Norman, 49).

Scotland: Salisbury Crags; Edinburgh; Lanarkshire; Rothesay; (Scott, 68): Killwinning; (John Smith *vide* Robertson, 57): Highgate; (Bate and Westwood, 1).

Ireland: Leixlip, Co. Dublin; Tempo, Co. Fermanagh; Goresbridge, Co. Kilkenny; (R.F.S.)

FOREIGN DISTRIBUTION:—

Europe: France; (25); Sweden; Norway; Denmark; Germany; Bohemia; Holland; Belgium; Turkey; Caucasus; (59).

North America; (59).

Family—ARMADILLIDIIDÆ.

Tail appendages not projecting when the animal is walking.

Genus—ARMADILLIDIUM Brandt, 1833 (3), p. 184.

Flagellum, with two joints; outer division of the tail appendages expanded and broader at the hinder end.

The members of the genus *Armadillidium* are more likely to be confounded, by the uninitiated, with the "Pill-millipedes" than with other Woodlice. Excepting *Cylisticus* (which has long pointed tail appendages) no other British forms have the power of rolling themselves up into a complete ball. The very arched body is characteristic of *Armadillidium*, and so is the groove into which the basal joints of the antennæ fit when the creatures curl up.

The first two abdominal appendages only are provided with air-tubes.

Armadillidium nasatum Budde-Lund. [PLATE XXII.](#)

1885 *Armadillidium nasatum* Budde-Lund (8), p. 51.

1892 *Armadillidium nasatum* Dollfus (14), p. 10, fig. 12.

1899 *Armadillidium nasatum* Norman (misprinted *Porcellidium*) (49), p. 57. pl. VI., figs. 5-8.

Armadillidium nasatum has a narrow but very prominent frontal lobe, which is almost square and curves somewhat upwards and backwards. The joints of the flagellum are approximately equal, and are together of the same length as the last peduncular joint.




FIG. 56.—FLAGELLUM
AND LAST PEDUNCULAR
JOINT OF THE ANTENNA
OF *Armadillidium
nasatum*.

The telson is as long as it is broad at the base, and tapers to a roundish point, while its sides are slightly incurved.

The outer divisions of the tail appendages are considerably longer than broad, and are more or less paddle-shaped.

It will be noticed that the slope from thorax to telson is more gentle than in the common species, *Armadillidium vulgare*, and the first thoracic segment is not so greatly developed. Consequently the species which we are considering does not produce a perfect sphere, and the antennæ are not hidden when it rolls up. It is interesting to compare this species with *Cylisticus convexus*. The surface of the body is smooth, and its colour is a delicate brownish grey with more or less distinct rows of darker markings.

[Pg 41]

BRITISH LOCALITIES:—

England: Maldon; (W.M.W. from R.M.): Bluebell Hill, Maidstone (W.M.W.); Clifton, banks of the Avon; (W.M.W. from J.T.C. 1900): Leigh Woods, Clifton; Tunbridge Wells; South Devon; (Stebbing in 49); Cheddar Cliffs, Somerset; (Norman, 49).

FOREIGN DISTRIBUTION:—

Europe: France; (28); Spain; (12); Italy; (23).

Armadillidium vulgare Latreille. [PLATE XXIII.](#)

- 1804 *Armadillo vulgaris* Latreille (37), p. 48.
- 1804 *Armadillo variegatus* Latreille (37).
- 1853 *Armadillo ater* Schnitzler (65), p. 48.
- 1816 *Armadillo maculatus* Risso (56), p. 158.
- 1818 *Armadillo pillularis* Say (62), p. 432.
- 1825 *Armadillo pustulosus* Dermarest (11), p. 323, pl. XLIX.
- 1830-4 *Armadillidium commutatum* Brandt and Ratzeburg (4), p. 81, pl. XIII., fig. 123.
- 1833 *Armadillidium zenckeri* Brandt (3), p. 185.
- 1839 *Armadillo trivialis* Koch (34), part 28, pl. XIV.
- 1898 *Armadillidium vulgare* Sars (59), p. 189, pl. LXXXII.

The common pill woodlouse is *Armadillidium vulgare*. Its frontal lobe is not large, though it is broad, while its margin where it joins the head is rounded and slightly recurved. The proximal joint of the flagellum is somewhat the shorter and the two together, as in *Armadillidium nasatum*, are of about the same length as the last joint of the peduncle.

The telson has the form of a triangle with the angles truncated and is about as long as it is broad at the base. The outer divisions of the tail appendages are considerably broader than they are long.

The species can roll itself up into a very perfect sphere, and when it assumes this form its antennæ are hidden beneath the much expanded lateral plates of the first thoracic segment.

The body is smooth, shiny, and strongly arched. Its colour varies very considerably, generally it is of a slaty-grey, but yellow markings are often present to a greater or less extent. In a specimen before us (from Bluebell Hill, Maidstone) the head is of a uniform dark grey, the sides of the thorax are yellow, while the back is mottled




FIG. 57.—FLAGELLUM
AND LAST PEDUNCULAR
JOINT OF THE ANTENNA
OF *Armadillidium
vulgare*.

[Pg 42]

with the same colour. The abdomen, including the telson, is also yellow with the exception only of the tail appendages, which are dark grey.

BRITISH LOCALITIES:—

England: Warley; (W.M.W.): Maldon; (W.M.W. from R.M.): Brightlingsea; Hanwell; Mortlake; Bluebell Hill, Maidstone; Langley; Skirmett; Pamber Forest; Kingston-on-Soar; Ipswich; (W.M.W.); Lynmouth; (W.M.W. from J.T.C.)

Scotland: (Scott, 68.)

Ireland: Ardrahan; (Norman, 50): Borris, Co. Carlow; Glandare, Terneay, Co. Cork; Courtstown, Co. Wexford; Cappagh, Co. Waterford; Castel, Co. Tipperary; (R.F.S.)

FOREIGN DISTRIBUTION:—

Europe: Throughout; (23).

Asia: Damascus; (23).

Africa: Algeria; (23).

Atlantic Isles: (23).

America: North and South; (23).

Australia: Melbourne (64 quoting Budde-Lund); New Zealand; (23).

Armadillidium pulchellum Zencker.

[PLATE XXIV.](#)

1799 *Oniscus pulchellus* Zencker (78) (quoted by Koch in Panzer), part 62, pl. XXI.

1833 *Armadillidium pulchellum* Brandt (3), p. 188.

1861 *Armadillo maculatus* Sill (69), p. 5.

1870 *Armadillidium pictum* Plateau (not Brandt) (55), p. 116.

1898 *Armadillidium pulchellum* Sars (59), p. 191, pl. LXXXIII., fig. 4.




FIG. 58.—FLAGELLUM AND LAST PEDUNCULAR JOINT OF THE ANTENNA OF *Armadillidium pulchellum*.

The smallest British species is *Armadillidium pulchellum*. The frontal lobe projects so as to make the head somewhat triangular. The antennæ are very short and the distal joint of the flagellum is three times the length of the other, while the two together are not as

long as the last peduncular joint.

The telson is truncated at the end so that it is by no means as long as it is broad at the base, and the outer divisions of the tail appendages are in similar proportion.

The colour of the body (which is smooth) is dark brown with four important series of light patches running down the back and less marked variegations between them.

BRITISH LOCALITIES:—

England: Matlock; (T.R.R.S.) Arnside; Westmorland (Brady, 50a).

Ireland: Ballymote, (Irish Nat., May, 1901), Sligo; (Scharff).

FOREIGN DISTRIBUTION:—

Europe: Vosges; Switzerland; Pyrenees; (23): Forest de Soignes; Belgium; (14).

Armadillidium depressum Brandt. [PLATE XXV.](#)

1833 *Armadillidium depressum* Brandt (3), p. 82. pl. XII., figs, 4, 5, 6, C, D.

1892 *Armadillidium depressum* Dollfus (14), p. 17-18.

The frontal lobe in *Armadillidium depressum* is very prominent and much recurved. The antennæ are fairly long and while the two joints of the flagellum are nearly equal




FIG. 59.—FLAGELLUM
AND LAST PEDUNCULAR
JOINT OF THE ANTENNA
OF *Armadillidium
depressum*.

they are together not so long as the last peduncular joint.

The telson is slightly longer than it is broad at the base, and its sides are incurved. As in *Armadillidium vulgare* and *Armadillidium pulchellum* the outer divisions of the tail appendages are broader than they are long.

The body is flatter than in the other species and bears tubercles; its colour is a slate-grey with yellowish markings. The first thoracic segment is well developed and the head appears as if almost completely imbedded in it.

BRITISH LOCALITIES:—

England: Clifton, banks of the Avon; (W.M.W. from J.T.C., 1900); Shirehampton, near Bristol; (Stebbing in 49); Clifton; (Dollfus from Miers 14).

FOREIGN DISTRIBUTION:—

Europe: France; Italy; (25); Asia Minor; (14, quoting Brandt).

Distribution of Species. There are not sufficient records at present to enable us to draw any conclusions as to the general distribution of Woodlice in the British Isles, but it is hoped that more attention will be given to these creatures, and that before long there may be other material available.

Conclusion. At the beginning it was mentioned that the present work grew out of an investigation into the fauna of Essex, and in order to show what may be expected when places are explored in which no collecting has been done, we may briefly indicate the results which we obtained in the county in question.

It was not long before a species new to Britain—to wit, *Porcellio ratzeburgii*—was found (74). This discovery was mentioned by Mr. Stebbing in the *Victoria County History of Essex* (p. 71), and he prophesied that most of the British species then would be met with in the county. We may safely claim to have shown that his prediction was true, for we have been able to record in the preceding pages no less than sixteen other species, as will be seen from the following lists:—

WOODLICE RECORDED FROM ESSEX.

- | | |
|---------------------------------------|----------------------------------|
| 1. <i>Ligia oceanica</i> | 10. <i>Porcellio pictus</i> |
| 2. <i>Ligidium hypnorum</i> | 11. <i>Porcellio dilatatus</i> |
| 3. <i>Trichoniscus pusillus</i> | 12. <i>Porcellio laevis</i> |
| 4. <i>Trichoniscus roseus</i> | 13. <i>Porcellio ratzeburgii</i> |
| 5. <i>Haplophthalmus danicus</i> | 14. <i>Metoponorthus pruinus</i> |
| 6. <i>Oniscus asellus</i> | 15. <i>Cylisticus convexus</i> |
| 7. <i>Philoscia muscorum</i> | 16. <i>Armadillidium nasatum</i> |
| 8. <i>Platyarthrus hoffmannseggii</i> | 17. <i>Armadillidium vulgare</i> |
| 9. <i>Porcellio scaber</i> | |

Of these *Ligidium hypnorum* calls for special mention, as it had not been found in this country since Mr. Stebbing discovered it in Surrey in 1873. Several of the *Porcellios* and *Cylisticus convexus* have been met with in but few places, and the same may be said of *Armadillidium nasatum*.

Of the British species not as yet found in Essex *Trichoniscus vividus* has at present only been recorded from Ireland; *Philoscia couchii* and *Armadillidium depressum* have not been collected except in the extreme south west of England, while *Metoponorthus cingendus* has hitherto only been noticed in Devonshire and Ireland. The other four species, with the exception of *Porcellio rathkei*, which is well

distributed in west Middlesex (and might have been expected to occur in Essex), are still rare. In fact, for *Trichoniscoides albidus* but two British localities are known; for *Haplophthalmus mengii* three (two in England and one in Ireland); while *Armadillidium pulchellum* has only been recorded from two or three places.

In other counties quite as satisfactory results were obtained as in Essex—a systematic search in Buckinghamshire brought to light at Eton three species which at the time had not been recorded from the British Isles, while in Middlesex, no less than a dozen species were found at Hanwell.

[Pg 45]

Since part of this contribution was printed our attention has been drawn to some notes by the Rev. Canon Norman and Professor G. S. Brady (50a). These bear out the remarks which have already been made, for among the species found by Professor Brady in the north of England were *Trichoniscoides albidus*, *Haplophthalmus mengii*, *Porcellio rathkei*, and *Armadillidium pulchellum*. In one of Canon Norman's previous papers (50—1903) he claims to have added the second species to the British list, and in the notes in question a similar claim is made with regard to the first and third. It should, however, be pointed out that all three of them were found in Buckinghamshire in 1899 by Mr. Webb, and that they were exhibited at the Nature Study Exhibition held in London in August, 1902. A specimen of *Armadillidium pulchellum* from Matlock was sent to us by the Rev. T. R. R. Stebbing in January, 1904, and was found, we understand, some considerable time previously.

It only remains for us to express our hearty thanks to the numerous friends and correspondents who have given us their ready help. The names of these have been printed in the text, but we would like to mention more particularly Dr. Calman, of the British Museum (Natural History), Monsieur Adrian Dollfus, Mr. Roland Matthams, the Rev. Canon Norman, Dr. Scharff, the Rev. T. R. R. Stebbing, and Miss Willmott.

FOOTNOTES:

- [1] "The Non-Marine Molluscs of Essex," by Wilfred Mark Webb; ESSEX NATURALIST, Vol. x. (1897), pp. 27-48 and 65-81.
- [2] The numbers in brackets refer to papers mentioned in the Bibliography at the end.
- [3] It should be pointed out that the methylated spirit now sold in the shops contains mineral naphtha and goes milky on the addition of water. Permission can be obtained from Somerset House to buy what is still called "ordinary methylated spirit," but at present five gallons has to be purchased at one time.

BIBLIOGRAPHY.

Explanation of the initials given in the text:—

B.M.—Specimens in the British Museum (Natural History).	C.S.—Charles Sillem.
J.T.C.—John Thomas Carrington.	T.R.R.S.—The Rev. Thomas R. R. Stebbing.
J.A.M.—James A. Murie.	W.M.W.—Wilfred Mark Webb.
R.M.—Roland Matthams.	R.W.—R. Welch.
R.F.S.—R. F. Scharff.	

- (1) BATE, C. SPENCE, and WESTWOOD, J.C.: *A history of the British Sessile-eyed Crustacea*, London, 1868.
- (2) BOSC, L. A. G.: *Manuel de l'histoire Naturelle des Crustacés*, Vol. II. Paris, 1830.
- (3) BRANDT, J. F.: "Conspectus Monographiae Crustaceorum Oniscodorum Latreillei," *Bull. Soc. Nat., Moscow*. Vol. VI. (1833), pp. 171-193.
- (4) BRANDT, J. F., and RATZEBURG, J. T. C.: *Medizinische Zoologie*. Vol. II., Berlin, 1830-1834.

- (5) BRUNTZ, L.: "Contribution à l'Etude de l'Excretion chez les Arthropodes." *Archives de Biologie*. Vol. XXII. (Nov., 1903), pp. 215-422, pls. I-VIII.
- (6) BUDDE-LUND, G.: "Danmarks Isopode Landkrebssdyr," *Naturhistorisk Tidsskrift*. Part 3, vol. VII. (1870, pp. 217-245).
- (7) BUDDE-LUND, G.: *Prospectus generum specierumque Crustaceorum Isopodum terrestrium*. Copenhagen, 1879.
- (8) BUDDE-LUND, G.: *Crustacea Isopoda Terrestria*. Hauniae, 1885.
- (9) CUVIER, G.: "Memoires sur les Cloportes terrestres." *Journ. d'hist. nat.* Vol. II. 1792.
- (10) DE GEER, C.: *Memoires pour servir à l'histoire des insectes*. Vol. VII., Stockholm, 1778.
- (11) DESMAREST, A.G.: *Considérations générales sur la classe des Crustacés*. Paris, 1825.
- (12) DOLLFUS, A.: "Catalogue raisonné des Isopodes terrestres de l'Espagne." *Ann. Soc. Espan. Hist. Nat.* Vol. XXI. (1892), pp. 161-190.
- (13) DOLLFUS, A.: "Sur la Distribution Geographique des Isopodes terrestres dans la region des Basses-Pyrenees." *Assoc. Francaise pour l'avancement des Sciences. Rep. Congress de Pau.*, 1892.
- (14) DOLLFUS, A.: "Le Genre Armadillidium." *Feu. des Jeunes Naturalistes*, Ser. 3. May 1892.
- (15) DOLLFUS, A.: "Catalogue raisonné des Isopodes terrestres de l'Espagne." Premier supplement, *Ann. Soc. Espan. Hist. Nat.* Vol. XXII. (1893), pp. 47-51.
- (16) DOLLFUS, A.: "Isopodes Terrestres in 'Voyage de M. Ch. Allmand aux Iles Canaries.'" *Mem. Soc. Zool. de France*. Vol. VI. (1893), p. 46.
- (17) DOLLFUS, A.: "Sur la Distribution Geographique des Armadilliens en Europe." *Compte-Rendu troisieme Congres. Internal. de Zool.* Leyden, Sept. 1895 (1896), pp. 356-358.
- (18) DOLLFUS, A.: "Les Isopodes terrestres du Nord de l'Afrique du Cap. Blanc a Tripoli." *Mem. Soc. Zool. Trans.* Vol. IX. (1896), pp. 523-553.
- (19) DOLLFUS, A.: *Crustacés Isopodes de la Sicile*. 1896.
- (20) DOLLFUS, A.: "On West Indian Isopod Crustaceans." *Proc. Zool. Soc. Lond.* 1896, pp. 388-400.
- (21) DOLLFUS, A.: "Tableau Iconographique des *Philoscia* d'Europe." *Feu. des Jeun. Natural.* Ser. 3, 1897, pp. 70-73 and 91-95.
- (22) DOLLFUS, A.: "Land Isopods der Balkan region." *Wissenschaft Mittheilung aus Bosnien und der Hercegovina*. Vol. IV. (1896).
- (23) DOLLFUS, A.: "Les Crustacés Isopodes Terrestres à grande dispersion," *Feu des Jeun. Natural.* Ser. 3 (Oct. 1897).
- (24) DOLLFUS, A.: "Sur la Distribution Geographique des Isopodes Terrestres dans l'Afrique Septentrional du Senegal à Obock." *Proc. Inter. Congres. Zool.* Cambridge, 1898, pp. 249-259.
- (25) DOLLFUS, A.: "Catalogue des Crustacés Isopodes Terrestres de France." *Feu. des Jeun. Natural.* Ser. 3 (October 1899).
- (26) FABRICIUS, J.C.: *Entomologia systematica*. Vol. II., Hauniae, 1793.
- (27) FABRICIUS, J. C.: Supplement to the last, 1798.
- (28) FERNIE, Dr.: *Animal Simples*, 1899.
- (29) FITCH, Asa.: *Noxious Insects of New York*. Albany, 1856.
- (30) GUERIN, F.E.: "Sur *Porcellio Poeyi*." *Ann. Sci. Soc. Entom. de France*. Vol. VI. (Paris, 1837, in Bull. ent., page 6).
- (31) HELLER, C.: *Reise der Novara; Crustacea*. 1865.
- (32) KINAHAN, J.R.: "Analysis of certain genera of terrestrial Isopoda." *Nat. Hist. Rev.* Vol. IV. (1857).
- (33) KINAHAN, J.R.: "On the genera *Philoscia*, *Itea* and *Philougria*." *Nat. Hist. Rev.* Vol. V. 1858.
- (34) KOCH, C.L.: *Deutschlands Crustaceen, Myriapoden and Arachniden*. Regensburg, 1835-1844.
Koch's descriptions and figures were published in *Deutschlands Crustaceen* and appeared also in Panzer's *Faunæ Insectorum Germanicæ* continued by Herrich-Schäffer (which see). The number of the parts of the latter which correspond to those of the former are placed in square brackets after them in the list here given. Part 6 (1836) [139]; Part 22 (1838) [162]; Part 28 (1839) [178]; Part 34 (1840) [180]; Part 36 (1844) [186]. Some of the dates quoted will be found to differ from those usually given (in Budde-Lund (8, p. 7) for instance), and printed on the parts of the copy in the Zoological Society's Library. Our authority for this change is Mr. C. D. Sherborne, who bases his dates upon reviews which he has discovered.
- (35) KOCH, C. L.: *System der Myriapoden mit den Verzeichnissen und Berichtigungen zu Deutschlands Crustaceen, Myriapoden und Arachniden*. Regensburg, 1847.
- (36) LAMARCK, J. B.: *Histoire naturelle des animaux sans vertébrés*. Vol. V., 1818.
- (37) LATREILLE, P. A.: *Histoire Naturelle des Crustacés et des Insectes*. Vol. VII., Paris, 1804.
- (38) LEACH, W. E.: "Tabular view of the external characters of four classes of animals, which Linné arranged under Insectes." *Trans. Linn. Soc. Lond.* Vol. XI., 1815.
- (39) LEREBOLLETT, A.: "Memoire sur les Crustacés de la famille des Cloportides qui habitent les environs de Strasbourg." *Mem. Soc. Mus. Hist. Nat. Strasbourg*. Vol. IV., 1853, pp. 130, pl. X.
- (40) LEUCKART, R.: *Verzeichnisz der zur Fauna Helgolands gehörenden wirbellosen Seethiere*. Braunsweig, 1847.

- (41) LINNÉ, C.: *Fauna Suecica*. First edition, Holmiæ, 1746.
- (42) LINNÉ, C.: *Fauna Suecica*. Second edition, Holmiæ, 1761.
- (43) LINNÉ, C.: *Systema Naturae*, ed. 12, 1767.
- (44) MIERS, E. J.: "Catalogue of New Zealand Crustacea." *Ann. Mag. Nat. Hist., Ser. 4.*, Vol. XVII., 1876.
- (45) MIERS, E. J.: "On a collection of Crustacea (Decapoda and Isopoda), chiefly from South America." *Proc. Zool. Soc.*, 1877, p. 653-678.
- (46) MILNE-EDWARDS, H.: *Histoire naturelle des Crustacés*. Vol. III. Paris, 1840.
- (47) NICHOLSON and LYDDEKER: *Manual of Palæontology*, 1889, Vol. II., p. 559.
- (48) NORMAN, A. M.: "Note on the discovery of *Ligidium agile*, Persoon (*Zia saundersii* Stebbing)." *Ann. Mag. Nat. Hist. Ser. 4*, Vol. XI., p. 419.
- (49) NORMAN, A. M.: "British Land Isopoda." *Ann. Mag. Nat. Hist. (Ser. 7)*, Vol. III. (1899), pp. 71-78, pl. VI.
- (50) NORMAN, A. M.: Continuation of the last. Vol. XI., 1903, pp. 309-372.
- (50a) NORMAN, A.M.: and BRADY C.S.: "British Land Isopoda." Second Supplement *Ann. Mag. Nat. Hist. (Ser. 7)*, Vol. XIV. (1904), pp. 449-450.
- (51) PANZER, G. W. F.: *Faunæ Insectorum Germanicæ initia, oder Deutschlands Insecten*. Parts 1-110, Nurnberg, 1793-1813. (See Koch.)
- (52) PANZER, G. W. F.: Continuation by Herrich-Schäffer, Parts 111-190, 1829-1844. (See Koch.)
- (53) PARFITT, E.: "The Fauna of Devon." Sessile-Eyed Crustacea. *Trans. Devon Assoc. Sci.*, Sept. 1873.
- (54) PERSOON: No paper upon Woodlice by this author is known, but he is quoted in Panzer's *Deutschlands Insecten* (51), which see.
- (55) PLATEAU, F.: "Crustacés Isopodes Terrestres." *Bull. Acad. Roy. de Belgique, Ser. 2.*, Vol. XXIX., No. 2 (1870), p. 112.
- (56) RISSO, A.: *Histoire Naturelle des Crustacés des environs de Nice*. Paris, 1816.
- (57) ROBERTSON, D.: "Catalogue of the Amphipoda and Isopoda of the Firth of Clyde." *Trans. Nat. Hist. Soc. Glasgow*, Vol. II. (1888), pp. 9-99.
- (58) ROULE, LOUIS: "Etudes sur le Development de Crustacés." *Ann. Sci. Nat.*, Vol. XVIII. (1895), pp. 1-156, pls. I-X.
- (59) SARS, G. O.: *An account of the Crustacea of Norway*. Vol. II., 1896-1899.
- (60) SAUSSURE, H. de: "Diagnoses de quelques crustacés nouveaux des Antilles et du Mexique." *Rev. et Mag. de Zool. (Ser. 2)*, Vol. IX. (1857), pp. 304-308.
- (61) SAVIGNY, J. C., and AUDOUIN, V.: *Description de l'Egypte*. Vol. XXII., 1827.
- (62) SAY, T.: "An account of the Crustacea of the United States." *J. Acad. Nat. Sci. Philadelphia*, Vol. I., pt. II., 1818.
- (63) SCHARFF, R. F.: "The Irish Woodlice." *Irish Naturalist*, Vol. III., 1894, pp. 4-7 and 25-29, pl. II.
- (64) SCHARFF, R. F.: "The Woodlice of Co. Carlow." *Irish Naturalist*, 1895, p. 319.
- (65) SCHNITZLER, H. J.: *De Oniscineis agri Bonnensis*. (Thesis.), Cologne, 1853.
- (66) SCHOBL, J.: "*Typhloniscus*, eine neue blinde Gattung der Crustacea Isopoda." *Sitzungsberichte der math. naturw. Acad. Wiss. Wien*. Vol. XL. (1860), pp. 279-330.
- (67) SCOPOLI, J. A.: *Entomologia Carniolica*. Vindibonæ, 1763.
- (68) SCOTT, T.: "The Land and Freshwater Crustacea of the District around Edinburgh." *Proc. R. Phys. S. Edin.*, Vol. XI. (1890-91), p. 75.
- (68a) SCOTT, T.: *British Association Hand Book on the Natural History of Glasgow*. 1901; *Isopoda*, pp. 335 and 336.
- (69) SILL, VICTOR: "Beitrag zur Kenntniss der Crustaceen, Arachniden and Myriapoden Siebenburgens." *Verhandl. u. Mittheil. des Siebenburgischen Ver. für Naturwiss. zu Hermannstadt*, Vol. XII. (1861), p. 1-11.
- (70) STEBBING, T. R. R.: "On a Crustacean of the Genus *Zia*." *Ann. Mag. Nat. Hist. Ser. 4.*, Vol. XI. (1873).
- (71) STEBBING, T. R. R.: *The Victoria County History of Essex* (1903); Crustacea, pp. 27-28.
- (71a) STEBBING, T. R. R.: *The Victoria County History of Buckinghamshire* (1905); Crustacea.
- (72) VEJDOVSKYF: "Zur Morphologie der Antennen und Schalendruse der Crustaceen." *Zeit. Wiss. Zool.*, Vol. LXIX., p. 378.
- (73) VERHOEF, K. W.: "Ueber Palæarktische Isopoden." *Zool. Anz.*, Vol. XXIV.
- (74) WEBB, WILFRED MARK: "The occurrence in Essex of a species of Woodlouse (Isopoda) new to Britain (*Porcellio ratzeburgii*, Brandt)." *Essex Naturalist*, Vol. XI. (1899), p. 127.
- (75) WEBB, WILFRED MARK: "Notes on Woodlice" (Including extracts from a paper read before the North London Natural History Society by James B. Casserley). *Science Gossip*, Vol. VI., New Series (1900), pp. 295-296.
- (76) WEBER, MAX.: "Uber einige neue Isopoden der Niederlandischen

- (77) ZADDACH, E. G.: *Synopses crustaceorum Prussicorum prodromus Regiomonti*, 1844.
- (78) ZENKER, C. D.: No paper upon woodlice by this author is known, but he is quoted in Panzer's *Deutschlands Insecten* (51), which see.
- (79) ZITTEL, K. A. von: *Textbook of Palæontology*. English Translation 1900, p. 668.

INDEX.

- Abdomen, 2, [2](#)
Aegidae, 2
Air cavities, in abdominal appendages, 31
Air tubes, 1, 6, 31, 33, 34, 35, 36, 37, 39
albidus, *Trichoniscoides*, description of, 25
Alcohol, 16
Antennae, large, 2, [3](#)
 smaller, 2, [3](#)
Alimentary canal, 6, [7](#)
Ants, 31
 wood, 14
Anus, 31, inset facing [10](#)
Appendages, 2
 abdominal, 5, [6](#)
Archaeoniscus brodiei, 2
Armadillidæ, 17
 characters of, 18, 40
Armadillidium, 15, 38
 commutatum, 41
 depressum, 44
 description of, 43
 flagellum of, [43](#)
Armadillidium nasatum, 41, 44, Plate xxii.
 description of, 40
 flagellum of, [40](#)
Armadillidium pictum, 42
 pulchellum, 43, 44, 45, Plate xxxiv.
 description of, 42
 flagellum of, [42](#)
Armadillidium vulgare, 15, 40, 43, 44, Plate xxiii.
 description of, 41
 flagellum of, [41](#)
 moulting of, 12, 13
 zenckeri, 41
Armadillo ater, 41
 maculatus, 41, 42
 pillularis, 41
 pustulosus, 41
 trivialis, 41
 variegatus, 41
 vulgaris, 41
Arthropoda, 1
Arteries, 7, [7](#)
asellus, *Oniscus*, description of, 27
Avebury, Lord, 31

Baden, Miocene of, 2
Bate and Westwood, 17, 19, 30
Blastoderm, 10
Body cavity, 11, inset facing [10](#)
Brady, Professor C. S., 45
Branchial nephrocytes, 8
British Association, 19
Brood pouch, 4, [5](#), 9
Budde-Lund, 28

Calman, Dr., 45
Carpenter, 15
Cells, nucleated, 10, inset facing [10](#)
Cephalic nephrocytes, 8
Cheslip, 15
Chiselbob, 15

Church louse, 16
 Cicatricula, 10, inset facing [10](#)
cingendus, *Metoponorthus*, description of, 38
 Circulatory system, 6, 7
 Classification, 17
 Coal measures, 1
 Collection of Woodlice, methods of, 16
 Commissures, 8, [8](#)
 Conclusion, 43
convexus, *Cylisticus*, description of, 39
couchii, *Philoscia*, description of, 30
 Crabs, land, 1
 Crustacea, 1
 Cud worm, 15, 16
Cylisticus, 40
 characters of, 17, 18, 38
Cylisticus convexus, 44, Plate xxi.
 description of, 39
 flagellum of, [39](#)
 laevis, 35, 39
Cymothoa oceanica, 20

danicus, *Haplophthalmus*, description of, 27
depressum, *Armadillidium*, description of, 43
 Development, 9
 Digestive glands, 6, [7](#)
dilatatus, *Porcellio*, description of, 33
 Distribution of species, 43
 Dollfus, Adrian, 45

 Economic considerations, 12
 Egg, 9, 10, [10](#), [11](#), inset facing [10](#)
 segmentation of, 11, inset facing [10](#)
 Embryo, 10, [11](#)
 Endoderm, 10, inset facing [10](#)
 Essex, Woodlice recorded from, 44
 Excretory organs, 7
 Exhibition of species new to Britain, 45
 Exmoor, woodlice and ants on, 14
 External structure, 2

 Flagellum, 3, [3](#)
 Food, 13, 14

 Ganglia, cerebral 8
 Geological history, 1
 Genera, 17
 Gill, 5, [5](#)
 Gills, 7, [6](#)
Glomeris marginata, 15
 Grammar sows, 16
 Growth, time required for, 12
 Gullet, 6
 Gut, fore, inset facing [10](#)
 mid, inset facing [10](#)

 Habits, 12
 Hall, Miss Kate M., 31
Haplophthalmus, 22
 characters of, 17, 18, 26
 danicus, 44, Plate viii.
 description of, 27
 flagellum of, [27](#)
 elegans, 26
 mengii, 44, 45, Plate vii.
 description of, 26
 flagellum of, [26](#)

 Head, 2, [2](#), [3](#)
 Heart, 6, 7, [7](#), 11
 Herefordshire, Old Red Sandstone, 1
 History, geological, 1
hoffmannseggii, *Platyarthrus*, description of, 30
 Hog-louse, 15
 Hogs, little grey, 16

Hydro-cyanic acid gas for destroying woodlice, 14

hypnorum, *Ligidium*, description of, 21

Insects, 1

Intestine, 6, [7](#)

formation of, in embryo, 10, 11

Isle of Wight, Oligocene of, 2, Oolite of, 2

Isopoda, 1, 19

Itea crassicornis, 30

laevis, 22

mengii, 26

rosea, 24

riparia, 22

vivida, 23

Jurassic, Upper, 2

Kinahan, Professor, 19, 28, 30

Kitchenball, 15

Labels, 16

laevis, *Porcellio*, description of, 35

Ligia, 21

branchial nephrocytes, 8

characters of, 17, 18, 19, 20

hypnorum, 21

melanocephala, 29

oceanica, 44, Plate i.

description of, 20

excretory organs of, 7, 8

flagellum of, 20, [20](#)

scopulorum, 20

Ligiidae, 17

characters of, 19

Ligiidae, characters of, 17, 18, 19

Ligidium, 19

characters of, 17, 18, 21

hypnorum, 44, Plate ii.

description of, 21

excretory organs of, 7

in Surrey, 19

persoonii, 21

Lip, the lower, 4, [5](#)

the upper, 4, [5](#)

Lucre pig, 15

Lugdor, 15

Mandibles, 3, [4](#)

Manure, from stables, favourable to woodlice, 14

Maxillae, first, 3, [4](#)

second, 3, [4](#)

openings of excretory organs on second, 7

Maxillipeds, 3, [4](#)

mengii, *Haplophthalmus*, description of, 26

Mesoderm, 10, inset facing [10](#)

Methylated spirit, 16

Metoponorthus, 21

characters of, 18, 37

cingendus, description of, 38, Plate xx.

flagellum of, [38](#)

pruinus, 44, Plate xix.

description of, [37](#)

flagellum of, 37

simplex, 38

Miocene, 2

Middlesex, West, 35

Millipede, pill, 15, 40

Moult, 11

Moulting, process of, 12, 13, 14

Mouth appendages, 3, [4](#)

muscorum, *Philoscia*, description of, 29

Names, local, 15

nasatum, *Armadillidium*, description of, 40

Nephrocytes, branchial, 8

cephalic, 8
 Nerve cord, 8, [8](#)
 Nervous system, 8
 Norman, Rev. Canon A. M., 25, 27, 45

oceanica, Ligia, description of, 20
 Oenigen, Miocene of, 2
 Old-sows, 16
 Old women of the wood, little, 16
 Oligocene, 2
 Onisci, 17
 characters of, 27
 Oniscidae, 19
Oniscus, 28
 branchial nephrocytes, 8
 characters of, 17, 18, 27
 agilis, 21
 asellus, 4, 15, 28, 44, Plate ix.
Oniscus asellus, Albino in Epping Forest, 28
 description of, 27
 excretory organs of, 8
 flagellum of, [28](#)
 young examples of, 28
 convexus, 39
 fossor, 19, 28
 granulatus, 32
 murarius, 27
 muscorum, 29
 oceanicus, 20
 pulchellum, 42
 sylvestris, 29
 Oolite, 2
 Ovaries, 8, [8](#)
 Oviducts, 8, [8](#)

 Palmer, 16
 Penis, 9, [9](#)
 Peduncle, 3, [3](#)
Philoscia, 21, 28
 characters of, 17, 18, 29
 couchii, 44, Plate xi.
 description of, 30
 flagellum of, [30](#)
 marmorata, 29
 muscorum, 21, 44, Plate x.
 description of, 20
 flagellum of, [29](#)
Philougria celer, 29
 riparia, 22
 rosea, 24
 vivida, 23
 Physicians, college of, rules for preparing woodlice for medicinal use, 18
pictus, Porcellio, description of, 33
 Pill millipede, 15, 40
Platyarthrus, 22
 characters of, 17, 18, 39
 hoffmannseggii, 44, Plate xii.
 description of, 30
 flagellum of, [31](#)
Porcellio, 6, 28, 38
 characters of, 7, 18, 32
 armadilloides, 39
 asper, 32
 aztecus, 35
 brandtii, 32
 cinerascens, 35
 cingendus, 38
 cotillae, 35
 cubensis, 35
 degeerii, 35
 dilatatus, 34, 44, Plate xv.
 description of, 33
 flagellum of, [34](#)

dubius, 32, 33
ferrugineus, 34
flavipes, 35
flavo-vittatus, 37
frontalis, 37
graniger, 32
laevis, 39, 44, Plate xvii.
 description of, 35
 flagellum of, [35](#)
lugubris, 36
maculicornis, 37
melanocephalus, 33
mexicanus, 35
mixtus, 33
montezumae, 32
musculus, 35
nemorensis, 36
nigra, 32
paulensis, 32
pictus, 34, 44, Plate xiv.
 description of, 33
 flagellum of, [33](#)
poeyi, 35
pruinus, 37
quercum, 36
rathkei, 35, 36, 44, 45
 description of, 34, Plate xvi.
 flagellum of, [35](#)
ratzeburgii, 44, Plate xviii.
 description of, 36
 flagellum of, [36](#)
scaber, 28, 44, Plate xiii.
 description of, 32
 flagellum of, 32
 Milne Edwards, 33
spinifrons, 39
striatus, 34
sumichtasti, 35
syriacus, 35
tetramoerus, 34
trilineatus, 34
trivittatus, 34
truncatus, 37
urbicus, 35
Porcellionides, Miers, 37
 Preservation of Woodlice, methods of, 16
 Pro-ectoderm, 10, inset facing [10](#)
 Pro-endoderm, 10, inset facing [10](#)
 Protoplasm, formative, 10
pruinus, *Metoponorthus*, description of, 37
 pseudotracheae, 31
pulchellum, *Armadillidium*, description of, 42
pusillus, *Trichoniscus*, description of, 22
 Purbeck Beds, 2

 Quay-louse, 20
 Quay lowders, 20

rathkei, *Porcellio*, description of, 34
ratzeburgii, *Porcellio*, description of, 36
 Reproductive organs, 8, [8](#), [9](#)
roseus, *Trichoniscus*, description of, 24
 Roule, Professor Louis, 9

 Sandstone, old red, 1
 Sars, Professor G. O., 17, 23
scaber, *Porcellio*, description of, 32
 Scharff, Dr. R. F., 17, 19, 23, 45
 Seminal reservoir, 9, [9](#)
 Socchetre, 16
 Sow bug, 15
 Species, distribution of, 43
 St. Anthony's hogs, 16
 Stebbing, the Rev. T. R. R., 19, 21, 35, 44, 45
 Swanley Horticultural College, 14

Tertiary deposits, 2
 Testes, 9, [9](#)
 Thorax, 2, 3
 Thrush louse, 15
 Tiggyhog, 15
 Tracheae, 6, 31
 Trichoniscidæ, characters of, 18, 22
Trichoniscoides, 21, 22, 26
 characters of, 17, 18, 25
 albidus, 44, 45
 description of, 25
 flagellum of, [25](#)
Trichoniscus, 21, 22, 26
 characters of, 17, 18, 22
 pusillus, 22, 23, 24, 25, 44, Plate iii.
 description of, 22
 flagellum of, [23](#)
 pygmaeus, description of, 23
 roseus, 23, 25, 44, Plate v.
 description of, 24
 flagellum of, [24](#)
 vividus, 22, 25, 44, Plate iv.
 description of, 23
 flagellum of, [24](#)
 Tubes for specimens, 16
Typhloniscus steinii, 30

vividus, *Trichoniscus*, description of, 23
vulgare, *Armadillidium*, description of, 41

 Walking legs, 4, 5
 Webb, Mr. Wilfred Mark, 20, 25, 26, 35, 36, 45
 Welsh names for Woodlice, 16
 Westwood, Bate and, 17, 19, 30
 White arsenic for destroying Woodlice, 14
 Willmott, Miss, 45
 Wood-ants, 15
 Woodlice, in captivity, 12
 methods for getting rid of, 24
 position of, 1
 storage of specimens, 16
 Welsh names for, 16
 Woodlouse, common, 4

 Yolk, food, 10, 11, inset facing [10](#)

Zia agilis, 21 *melanocephala*, 29


LIST OF PLATES.

	PLATE
<i>Ligia oceanica</i> Linné	I.
<i>Ligidium hypnorum</i> Cuvier	II.
<i>Trichoniscus pusillus</i> Brandt	III.
<i>Trichoniscus vividus</i> Koch	IV.
<i>Trichoniscus roseus</i> Koch	V.
<i>Trichoniscoides albidus</i> Budde-Lund	VI.
<i>Haplophthalmus mengii</i> Zaddach	VII.
<i>Haplophthalmus danicus</i> Budde-Lund	VIII.
<i>Oniscus asellus</i> Linné	IX.
<i>Philoscia muscorum</i> Scopoli	X.
<i>Philoscia couchii</i> Kinahan	XI.
<i>Platyarthus hoffmannseggii</i> Brandt	XII.
<i>Porcellio scabier</i> Latreille	XIII.
<i>Porcellio pictus</i> Brandt and Ratzeburg	XIV.
<i>Porcellio dilatatus</i> Brandt	XV.
<i>Porcellio rathkei</i> Brandt	XVI.
<i>Porcellio laevis</i> Latreille	XVII.
<i>Porcellio ratzeburgii</i> Brandt	XVIII.
<i>Metoponorthus pruinosus</i> Brandt	XIX.
<i>Metoponorthus cingendus</i> Kinahan	XX.
<i>Cylisticus convexus</i> De Geer	XXI.
<i>Armadillidium nasatum</i> Budde-Lund	XXII.

BRITISH WOODLICE.

PLATE I.

[Plate i]



LIGIA OCEANICA Linné. THE QUAY-LOUSE.
Length, two to three centimetres.


Charles Sillem, del. ad nat.

F. W. Reader, sculpt.

BRITISH WOODLICE.

PLATE II.

[Plate ii]



LIGIDIUM HYPNORUM Cuvier.
Length, nine millimetres.


Charles Sillem, del. ad nat.

F. W. Reader, sculpt.

BRITISH WOODLICE.

PLATE III.

[Plate iii]



TRICHONISCUS PUSILLUS Brandt.
Length, four millimetres.


Charles Sillem, del. ad nat.

F. W. Reader, sculpt.

BRITISH WOODLICE.

PLATE IV.

[Plate iv]



TRICHONISCUS VIVIDUS Koch.
Length, eight millimetres.


Charles Sillem, del. ad nat.

F. W. Reader, sculpt.

BRITISH WOODLICE.

PLATE V.

[Plate v]



TRICHONISCUS ROSEUS Koch.
Length, five millimetres.


Charles Sillem, del. ad nat.

F. W. Reader, sculpt.

BRITISH WOODLICE.

PLATE VI.

[Plate vi]



TRICHONISCOIDES ALBIDUS Budde-Lund.
Length, four millimetres.


Charles Sillem, del. ad nat.

F. W. Reader, sculpt.

BRITISH WOODLICE.

PLATE VII.

[Plate vii]



HAPLOPHTHALMUS MENGII Zaddach.
Length, three to four millimetres.


Charles Sillem, del. ad nat.

F. W. Reader, sculpt.

BRITISH WOODLICE.

PLATE VIII.

[Plate
viii]



HAPLOPHTHALMUS DANICUS Budde-Lund.
Length, three to four millimetres.


Charles Sillem, del. ad nat.

F. W. Reader, sculpt.

BRITISH WOODLICE.

PLATE IX.

[Plate ix]



ONISCUS ASELLUS Linné (The common slater).

Length, sixteen millimetres.


Charles Sillem, del. ad nat.

F. W. Reader, sculpt.

BRITISH WOODLICE.

PLATE X.

[Plate x]



PHILOSCIA MUSCORUM Scopoli.

Length, nine millimetres.


Charles Sillem, del. ad nat.

F. W. Reader, sculpt.

BRITISH WOODLICE.

PLATE XI.

[Plate xi]



PHILOSCIA COUCHII Kinahan.
Length, nine millimetres.


Charles Sillem, del. ad nat.

F. W. Reader, sculpt.

BRITISH WOODLICE.

PLATE XII.

[Plate xii]



PLATYARTHUS HOFFMANNSEGGII Brandt.
Length, three millimetres.


Charles Sillem, del. ad nat.

F. W. Reader, sculpt.

BRITISH WOODLICE.

PLATE XIII.

[Plate
xiii]



PORCELLIO SCABER Latreille.
Length, fourteen millimetres.


Charles Sillem, del. ad nat.

F. W. Reader, sculpt.

BRITISH WOODLICE.

PLATE XIV.

[Plate
xiv]



PORCELLIO PICTUS Brandt and Ratzeburg.
Length, thirteen millimetres.


Charles Sillem, del. ad nat.

F. W. Reader, sculpt.

BRITISH WOODLICE.

PLATE XV.

[Plate xv]



PORCELLIO DILATATUS Brandt.
Length, fifteen millimetres.


Charles Sillem, del. ad nat.

F. W. Reader, sculpt.

BRITISH WOODLICE.

PLATE XVI.

[Plate
xvi]



PORCELLIO RATHKEI Brandt.
Length, twelve millimetres.


Charles Sillem, del. ad nat.

F. W. Reader, sculpt.

BRITISH WOODLICE.

PLATE XVII.

[Plate
xvii]



PORCELLIO LAEVIS Latreille.
Length, sixteen millimetres.


Charles Sillem, del. ad nat.

F. W. Reader, sculpt.

BRITISH WOODLICE.

PLATE XVIII.

[Plate
xviii]



PORCELLIO RATZEBURGHII Brandt.
Length, eleven millimetres.


Charles Sillem, del. ad nat.

F. W. Reader, sculpt.

BRITISH WOODLICE.

PLATE XIX.

[Plate
xix]



METOPONORTHUS PRUNOSUS Brandt.
Length, nine millimetres.


Charles Sillem, del. ad nat.

F. W. Reader, sculpt.

BRITISH WOODLICE.

PLATE XX.

[Plate xx]



METOPONORTHUS CINGENDUS Kinahan.
Length, six millimetres.


Charles Sillem, del. ad nat.

F. W. Reader, sculpt.

BRITISH WOODLICE.

PLATE XXI.

[Plate
xxi]



CYLISTICUS CONVEXUS De Geer.
Length, twelve millimetres.


Charles Sillem, del. ad nat.

F. W. Reader, sculpt.

BRITISH WOODLICE.

PLATE XXII.

[Plate
xxii]



ARMADILLIDIUM NASATUM Budde-Lund.
Length, fifteen millimetres.


Charles Sillem, del. ad nat.

F. W. Reader, sculpt.

BRITISH WOODLICE.

PLATE XXIII.

[Plate
xxiii]



ARMADILLIDIUM VULGARE Latreille.
Length, fifteen millimetres.


Charles Sillem, del. ad nat.

F. W. Reader, sculpt.

BRITISH WOODLICE.

PLATE XXIV.

[Plate
xxiv]



ARMADILLIDIUM PULCHELLUM Zencker.
Length, five millimetres.


Charles Sillem, del. ad nat.

F. W. Reader, sculpt.

BRITISH WOODLICE.

PLATE XXV.

[Plate
xxv]



ARMADILLIDIUM DEPRESSUM Brandt.
Length, fifteen millimetres..

Charles Sillem, del. ad nat.

F. W. Reader, sculpt.

Misprints and misspellings noted or corrected in the text:

Page vii [CONTENTS]: "Family--Ligidæ " changed to "Family--Ligiidæ".

Page vii [CONTENTS]: "*Haplophthalmus mengii*" changed to "*Haplophthalmus mengii*".

Page viii [CONTENTS]: "*Armydillidium vulgare*" changed to "*Armadillidium vulgare*".

Page 2: "Aegidae is found" changed to "Aegidae which is found".

Page 16: "naptha" changed to "naphtha" in footnote.

Page 28: "Faroe Islands; Thornsharn;" changed to "Faroe Islands; Thorsharn;".

Page 29: "1833 *Philoscia marmorata* ...", changed to "1833 *Philoscia marmorata* ..."

Page 30: Missing page ref. at end of "1868 *Philoscia couchii* Bate and Westwood (1), p."

Page 33: Missing page ref. at end of "1868 *Porcellio pictus* Bate and Westwood (1), p."

Page 33: Missing page ref. at end of "1868 *Porcellio dilatatus* Bate and Westwood (1), p."

Page 51 [INDEX]: "Ligidæ, characters of" changed to "Ligiidæ, characters of".

[Back to top](#)

*** END OF THE PROJECT GUTENBERG EBOOK THE
BRITISH WOODLICE ***

Updated editions will replace the previous one—the old editions will be renamed.

Creating the works from print editions not protected by U.S. copyright law means that no one owns a United States copyright in these works, so the Foundation (and you!) can copy and distribute it in the United States without permission and without paying copyright royalties. Special rules, set forth in the General Terms of

Use part of this license, apply to copying and distributing Project Gutenberg™ electronic works to protect the PROJECT GUTENBERG™ concept and trademark. Project Gutenberg is a registered trademark, and may not be used if you charge for an eBook, except by following the terms of the trademark license, including paying royalties for use of the Project Gutenberg trademark. If you do not charge anything for copies of this eBook, complying with the trademark license is very easy. You may use this eBook for nearly any purpose such as creation of derivative works, reports, performances and research. Project Gutenberg eBooks may be modified and printed and given away—you may do practically ANYTHING in the United States with eBooks not protected by U.S. copyright law. Redistribution is subject to the trademark license, especially commercial redistribution.

START: FULL LICENSE
THE FULL PROJECT GUTENBERG LICENSE
PLEASE READ THIS BEFORE YOU DISTRIBUTE OR USE THIS
WORK

To protect the Project Gutenberg™ mission of promoting the free distribution of electronic works, by using or distributing this work (or any other work associated in any way with the phrase “Project Gutenberg”), you agree to comply with all the terms of the Full Project Gutenberg™ License available with this file or online at www.gutenberg.org/license.

**Section 1. General Terms of Use and
Redistributing Project Gutenberg™
electronic works**

1.A. By reading or using any part of this Project Gutenberg™ electronic work, you indicate that you have read, understand, agree to and accept all the terms of this license and intellectual property (trademark/copyright) agreement. If you do not agree to abide by all the terms of this agreement, you must cease using and return or destroy all copies of Project Gutenberg™ electronic works in your possession. If you paid a fee for obtaining a copy of or access to a Project Gutenberg™ electronic work and you do not agree to be bound by the terms of this agreement, you may obtain a refund from the person or entity to whom you paid the fee as set forth in paragraph 1.E.8.

1.B. “Project Gutenberg” is a registered trademark. It may only be used on or associated in any way with an electronic work by people who agree to be bound by the terms of this agreement. There are a few things that you can do with most Project Gutenberg™ electronic works even without complying with the full terms of this agreement. See paragraph 1.C below. There are a lot of things you can do with Project Gutenberg™ electronic works if you follow the terms of this agreement and help preserve free future access to Project Gutenberg™ electronic works. See paragraph 1.E below.

1.C. The Project Gutenberg Literary Archive Foundation (“the Foundation” or PGLAF), owns a compilation copyright in the collection of Project Gutenberg™ electronic works. Nearly all the individual works in the collection are in the public domain in the United States. If an individual work is unprotected by copyright law in the United States and you are located in the United States, we do not claim a right to prevent you from copying, distributing, performing, displaying or creating derivative works based on the work as long as all references to Project Gutenberg are removed. Of course, we hope that you will support the Project Gutenberg™ mission of promoting free access to electronic works by freely sharing Project Gutenberg™ works in compliance

with the terms of this agreement for keeping the Project Gutenberg™ name associated with the work. You can easily comply with the terms of this agreement by keeping this work in the same format with its attached full Project Gutenberg™ License when you share it without charge with others.

1.D. The copyright laws of the place where you are located also govern what you can do with this work. Copyright laws in most countries are in a constant state of change. If you are outside the United States, check the laws of your country in addition to the terms of this agreement before downloading, copying, displaying, performing, distributing or creating derivative works based on this work or any other Project Gutenberg™ work. The Foundation makes no representations concerning the copyright status of any work in any country other than the United States.

1.E. Unless you have removed all references to Project Gutenberg:

1.E.1. The following sentence, with active links to, or other immediate access to, the full Project Gutenberg™ License must appear prominently whenever any copy of a Project Gutenberg™ work (any work on which the phrase “Project Gutenberg” appears, or with which the phrase “Project Gutenberg” is associated) is accessed, displayed, performed, viewed, copied or distributed:

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

1.E.2. If an individual Project Gutenberg™ electronic work is derived from texts not protected by U.S. copyright law (does not contain a notice indicating that it is posted with permission of the copyright holder), the work can be copied and distributed to anyone in the United States without paying any fees or charges. If you are redistributing or providing access to a work with the phrase “Project Gutenberg” associated with or appearing on the work, you must comply either with the requirements of paragraphs 1.E.1 through 1.E.7 or obtain permission for the use of the work and the Project Gutenberg™ trademark as set forth in paragraphs 1.E.8 or 1.E.9.

1.E.3. If an individual Project Gutenberg™ electronic work is posted with the permission of the copyright holder, your use and distribution must comply with both paragraphs 1.E.1 through 1.E.7 and any additional terms imposed by the copyright holder. Additional terms will be linked to the Project Gutenberg™ License for all works posted with the permission of the copyright holder found at the beginning of this work.

1.E.4. Do not unlink or detach or remove the full Project Gutenberg™ License terms from this work, or any files containing a part of this work or any other work associated with Project Gutenberg™.

1.E.5. Do not copy, display, perform, distribute or redistribute this electronic work, or any part of this electronic work, without prominently displaying the sentence set forth in paragraph 1.E.1 with active links or immediate access to the full terms of the Project Gutenberg™ License.

1.E.6. You may convert to and distribute this work in any binary, compressed, marked up, nonproprietary or proprietary form, including any word processing or hypertext form. However, if you provide access to or distribute copies of a Project Gutenberg™ work in a format other than “Plain Vanilla ASCII” or other format used in the official version posted on the official Project Gutenberg™ website (www.gutenberg.org), you must, at no additional cost, fee or expense to the user, provide a copy, a means of exporting a copy, or a means of obtaining a copy upon request, of the work in its original “Plain Vanilla ASCII” or other form. Any alternate format must include the full Project Gutenberg™ License as specified in paragraph 1.E.1.

1.E.7. Do not charge a fee for access to, viewing, displaying, performing, copying or distributing any Project Gutenberg™ works unless you comply with paragraph 1.E.8 or 1.E.9.

1.E.8. You may charge a reasonable fee for copies of or providing access to or distributing Project Gutenberg™ electronic works provided that:

- You pay a royalty fee of 20% of the gross profits you derive from the use of Project Gutenberg™ works calculated using the method you already use to calculate your applicable taxes. The fee is owed to the owner of the Project Gutenberg™ trademark, but he has agreed to donate royalties under this paragraph to the Project Gutenberg Literary Archive Foundation. Royalty payments must be paid within 60 days following each date on which you prepare (or are legally required to prepare) your periodic tax returns. Royalty payments should be clearly marked as such and sent to the Project Gutenberg Literary Archive Foundation at the address specified in Section 4, “Information about donations to the Project Gutenberg Literary Archive Foundation.”
- You provide a full refund of any money paid by a user who notifies you in writing (or by e-mail) within 30 days of receipt that s/he does not agree to the terms of the full Project Gutenberg™ License. You must require such a user to return or destroy all copies of the works possessed in a physical medium and discontinue all use of and all access to other copies of Project Gutenberg™ works.
- You provide, in accordance with paragraph 1.F.3, a full refund of any money paid for a work or a replacement copy, if a defect in the electronic work is discovered and reported to you within 90 days of receipt of the work.
- You comply with all other terms of this agreement for free distribution of Project Gutenberg™ works.

1.E.9. If you wish to charge a fee or distribute a Project Gutenberg™ electronic work or group of works on different terms than are set forth in this agreement, you must obtain permission in writing from the Project Gutenberg Literary Archive Foundation, the manager of the Project Gutenberg™ trademark. Contact the Foundation as set forth in Section 3 below.

1.F.

1.F.1. Project Gutenberg volunteers and employees expend considerable effort to identify, do copyright research on, transcribe and proofread works not protected by U.S. copyright law in creating the Project Gutenberg™ collection. Despite these efforts, Project Gutenberg™ electronic works, and the medium on which they may be stored, may contain “Defects,” such as, but not limited to, incomplete, inaccurate or corrupt data,

transcription errors, a copyright or other intellectual property infringement, a defective or damaged disk or other medium, a computer virus, or computer codes that damage or cannot be read by your equipment.

1.F.2. LIMITED WARRANTY, DISCLAIMER OF DAMAGES - Except for the "Right of Replacement or Refund" described in paragraph 1.F.3, the Project Gutenberg Literary Archive Foundation, the owner of the Project Gutenberg™ trademark, and any other party distributing a Project Gutenberg™ electronic work under this agreement, disclaim all liability to you for damages, costs and expenses, including legal fees. YOU AGREE THAT YOU HAVE NO REMEDIES FOR NEGLIGENCE, STRICT LIABILITY, BREACH OF WARRANTY OR BREACH OF CONTRACT EXCEPT THOSE PROVIDED IN PARAGRAPH 1.F.3. YOU AGREE THAT THE FOUNDATION, THE TRADEMARK OWNER, AND ANY DISTRIBUTOR UNDER THIS AGREEMENT WILL NOT BE LIABLE TO YOU FOR ACTUAL, DIRECT, INDIRECT, CONSEQUENTIAL, PUNITIVE OR INCIDENTAL DAMAGES EVEN IF YOU GIVE NOTICE OF THE POSSIBILITY OF SUCH DAMAGE.

1.F.3. LIMITED RIGHT OF REPLACEMENT OR REFUND - If you discover a defect in this electronic work within 90 days of receiving it, you can receive a refund of the money (if any) you paid for it by sending a written explanation to the person you received the work from. If you received the work on a physical medium, you must return the medium with your written explanation. The person or entity that provided you with the defective work may elect to provide a replacement copy in lieu of a refund. If you received the work electronically, the person or entity providing it to you may choose to give you a second opportunity to receive the work electronically in lieu of a refund. If the second copy is also defective, you may demand a refund in writing without further opportunities to fix the problem.

1.F.4. Except for the limited right of replacement or refund set forth in paragraph 1.F.3, this work is provided to you 'AS-IS', WITH NO OTHER WARRANTIES OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR ANY PURPOSE.

1.F.5. Some states do not allow disclaimers of certain implied warranties or the exclusion or limitation of certain types of damages. If any disclaimer or limitation set forth in this agreement violates the law of the state applicable to this agreement, the agreement shall be interpreted to make the maximum disclaimer or limitation permitted by the applicable state law. The invalidity or unenforceability of any provision of this agreement shall not void the remaining provisions.

1.F.6. INDEMNITY - You agree to indemnify and hold the Foundation, the trademark owner, any agent or employee of the Foundation, anyone providing copies of Project Gutenberg™ electronic works in accordance with this agreement, and any volunteers associated with the production, promotion and distribution of Project Gutenberg™ electronic works, harmless from all liability, costs and expenses, including legal fees, that arise directly or indirectly from any of the following which you do or cause to occur: (a) distribution of this or any Project Gutenberg™ work, (b) alteration, modification, or additions or deletions to any Project Gutenberg™ work, and (c) any Defect you cause.

Section 2. Information about the Mission of Project Gutenberg™

Project Gutenberg™ is synonymous with the free

distribution of electronic works in formats readable by the widest variety of computers including obsolete, old, middle-aged and new computers. It exists because of the efforts of hundreds of volunteers and donations from people in all walks of life.

Volunteers and financial support to provide volunteers with the assistance they need are critical to reaching Project Gutenberg™'s goals and ensuring that the Project Gutenberg™ collection will remain freely available for generations to come. In 2001, the Project Gutenberg Literary Archive Foundation was created to provide a secure and permanent future for Project Gutenberg™ and future generations. To learn more about the Project Gutenberg Literary Archive Foundation and how your efforts and donations can help, see Sections 3 and 4 and the Foundation information page at www.gutenberg.org.

Section 3. Information about the Project Gutenberg Literary Archive Foundation

The Project Gutenberg Literary Archive Foundation is a non-profit 501(c)(3) educational corporation organized under the laws of the state of Mississippi and granted tax exempt status by the Internal Revenue Service. The Foundation's EIN or federal tax identification number is 64-6221541. Contributions to the Project Gutenberg Literary Archive Foundation are tax deductible to the full extent permitted by U.S. federal laws and your state's laws.

The Foundation's business office is located at 809 North 1500 West, Salt Lake City, UT 84116, (801) 596-1887. Email contact links and up to date contact information can be found at the Foundation's website and official page at www.gutenberg.org/contact

Section 4. Information about Donations to the Project Gutenberg Literary Archive Foundation

Project Gutenberg™ depends upon and cannot survive without widespread public support and donations to carry out its mission of increasing the number of public domain and licensed works that can be freely distributed in machine-readable form accessible by the widest array of equipment including outdated equipment. Many small donations (\$1 to \$5,000) are particularly important to maintaining tax exempt status with the IRS.

The Foundation is committed to complying with the laws regulating charities and charitable donations in all 50 states of the United States. Compliance requirements are not uniform and it takes a considerable effort, much paperwork and many fees to meet and keep up with these requirements. We do not solicit donations in locations where we have not received written confirmation of compliance. To SEND DONATIONS or determine the status of compliance for any particular state visit www.gutenberg.org/donate.

While we cannot and do not solicit contributions from states where we have not met the solicitation requirements, we know of no prohibition against accepting unsolicited donations from donors in such states who approach us with offers to donate.

International donations are gratefully accepted, but we cannot make any statements concerning tax treatment of donations received from outside the United States. U.S. laws alone swamp our small staff.

Please check the Project Gutenberg web pages for current donation methods and addresses. Donations are

accepted in a number of other ways including checks, online payments and credit card donations. To donate, please visit: www.gutenberg.org/donate

Section 5. General Information About Project Gutenberg™ electronic works

Professor Michael S. Hart was the originator of the Project Gutenberg™ concept of a library of electronic works that could be freely shared with anyone. For forty years, he produced and distributed Project Gutenberg™ eBooks with only a loose network of volunteer support.

Project Gutenberg™ eBooks are often created from several printed editions, all of which are confirmed as not protected by copyright in the U.S. unless a copyright notice is included. Thus, we do not necessarily keep eBooks in compliance with any particular paper edition.

Most people start at our website which has the main PG search facility: www.gutenberg.org.

This website includes information about Project Gutenberg™, including how to make donations to the Project Gutenberg Literary Archive Foundation, how to help produce our new eBooks, and how to subscribe to our email newsletter to hear about new eBooks.